

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

Catalogue of Indian Insects

PART I—Acrydidae (Tettigidae)

BY

T. BAINBRIGGE FLETCHER,

R.N., F.L.S., F.E.S., F.Z.S.,

(Imperial Entomologist.)



CALCUTTA
SUPERINTENDENT GOVERNMENT PRINTING, INDIA
1921

Agents for the Sale of Books Published by the Superintendent of Government Printing, India, Calcutta

IN EUROPE

Constable & Co., 10, Orange Street, Leicester Square, London, W.C.
 Kegan Paul, Trench, Trübner & Co., 68-74, Carter Lane, E.C., and 39, New Oxford Street, London, W.C.
 Bernard Quaritch, 11, Grafton Street, New Bond Street, London, W.
 P. S. King & Sons, 2 & 4, Great Smith Street, Westminster, London, S.W.
 H. S. King & Co., 65, Cornhill, E.C., and 9, Pall Mall, London, W.
 Grindlay & Co., 54, Parliament Street, London, S.W.

Luzac & Co., 46, Great Russell Street, London, W.C.
 W. Thacker & Co., 2, Creed Lane, London, E.C.
 T. Fisher Unwin, Ltd., 1, Adelphi Terrace, London, W.C.
 Wheldon & Wesley, Ltd., 28, Essex Street, London, W.C.
 B. H. Blackwell, 50 & 51, Broad Street, Oxford.
 Deighton Bell & Co., Ltd., Cambridge.
 Oliver and Boyd, Tweeddale Court, Edinburgh.
 E. Ponsonby, Ltd., 116, Grafton Street, Dublin.
 Ernest Leroux, 28, Rue Bonaparte, Paris.
 Martinus Nijhoff, The Hague, Holland.

IN INDIA AND CEYLON

Thacker, Spink & Co., Calcutta and Simla.
 Newman & Co., Calcutta.
 B. Cambray & Co., Calcutta.
 S. K. Lahiri & Co., Calcutta.
 B. Banerjee & Co., Calcutta.
 The Indian School Supply Depôt, 309, Bow Bazar Street, Calcutta, and 226, Nawabpur, Dacca.
 Butterworth & Co. (India), Ltd., Calcutta.
 Rai M. C. Sarcar Bahadur and Sons, 90/2A, Harrison Road, Calcutta.
 The Weldon Library, 57, Park Street, West, Calcutta.
 Standard Literature Co., Limited, Calcutta.
 Lal Chand & Sons, Calcutta.
 Association Press, Calcutta.
 Higginbothams, & Co., Madras.
 V. Kalyanarama Iyer & Co., Madras.
 G. A. Natesan & Co., Madras.
 S. Murthy & Co., Madras.
 Thompson & Co., Madras.
 Temple & Co., Madras.
 P. R. Rama Iyer & Co., Madras.
 Vas & Co., Madras.
 E. M. Gopalakrishna Kone, Madura.
 Thacker & Co., Ltd., Bombay.
 D. B. Taraporevala, Sons & Co., Bombay.
 Mrs. Radhabai Atmaram Sagoon, Bombay.
 Sunder Pandurang, Bombay.
 Gopal Narayan & Co., Bombay.
 Ramchandra Govind & Son, Kalbadevi, Bombay.

Proprietor, New Kitabkhana, Poona.
 The Standard Bookstall, Karachi.
 Mangaldas Harkisandas, Surat.
 Karsandas Narandas & Sons, Surat.
 A. H. Wheeler & Co., Allahabad, Calcutta and Bombay.
 N. B. Mathur, Supt., Nazir Kanun-i-Hind Press, Allahabad.
 Munshi Seeta Ram, Managing Proprietor, Indian Army Book Depôt, Juhu, Cawnpore.
 Rai Sahib M. Gulab Singh & Sons, Mufid-i-Am Press, Lahore and Allahabad.
 Rama Krishna & Sons, Lahore.
 Supt., American Baptist Mission Press, Rangoon.
 Manager, the "Hitavada," Nagpur.
 S. C. Talukdar, Proprietor, Students and Company, Cooch Behar.
 A. M. & F. Ferguson, Colombo, Ceylon.
 Manager, Educational Book Depôts, Nagpur and Jubbulpore.*
 Manager of the Imperial Book Depôt, 63 Chandney Chauk Street, Delhi.*
 Manager, "The Agra Medical Hall and Co-operative Association, Ltd." (Successors to A. John & Co., Agra).*
 Supt., Basel Mission Book and Tract Depository, Mangalore.*
 P. Varadachary & Co., Madras.*
 Ram Dayal Agarwala, 184, Katra, Allahabad.*
 D. C. Anand & Sons, Peshawar.*
 Manager, Newal Kishore Press, Lucknow.*
 Maung Lu Gale, Proprietor, Law Book Depôt, Mandalay *

* Agents for the sale of Legislative Department publications only.

P R E F A C E .

THE Acrydidæ constitute a group of the short-horned grasshoppers which is considered by many authors to be a sub-family of the Acrididæ, from which, however, the Acrydidæ are distinguished by (1) the fact that the claws of the tarsi are not provided with an arolium (or pad between the two terminal claws; also known as a pulvillus) and (2) the prolongation of the pronotum backwards over the abdomen. The first of the two distinctions is constant and serves to mark the Acrydidæ sharply from the Acrididæ, but the backward projection of the pronotum is found in some Acridids also. Walker (*Ann. Ent. Soc. America* XII 285; Dec. 1919) finds grounds for considering that the Acrydidæ are more primitive than the Acrididæ. On the whole, therefore, it seems preferable to rank the Acrydids as a distinct family.

The names under which these small grasshoppers are best known to entomologists have usually been variants of the names *Tettix* and *Tetrix*, both of which are synonyms of *Acrydium*, and, as *Acrydium* is the earliest described genus in this group, the family name is derivable from that of this genus.

The Acrydidæ are small to moderately small insects, usually most abundant in damp grassy localities. Many species of the subfamily Scelimeninæ are sub-aquatic, never found far from water and usually occurring on wet rocks alongside streams and pools, diving into and swimming beneath the water when disturbed, the hind tibiæ in many species being expanded and flattened to form swimming legs.

Little is on record regarding the life-histories and habits of these insects. The immature stages are often found in company with the adults on damp rocks, etc., where both appear to feed on moss and other low forms of vegetable life. One species has been found as a minor pest of cabbages at Pusa.

The literature on this group is fairly extensive, the more important publications on the Indian forms being:—

- 1834. Westwood, *Zoological Journal*, Vol. V.
- 1839. Serville, *Hist. Nat. des Insectes*, Orthoptères.
- 1842. de Haan, in *Temminck's Verhandel.*, Orthoptera.
- 1861. de Saussure, "Etude sur quelques Orthoptères du musée de Genève" (*Ann. Soc. Ent. France* (4) I 469-494; 1860).
- 1871. Walker, *Cat. of Dermaptera Saltatoria*, Part V.

1871. Stål, Recensio Orthopterorum, Pars I.
1887. Bolivar, Essai sur les Acridiens de la tribu des Tettigidæ, (*Ann. Soc. Ent. Belg.* XXXI 175-313).
1893. Brunner, Révision du système des Orthopteres (*Ann. Mus. Genova* XXXIII 103-114, t. 5).
1898. Bolivar, Contributions à l'étude des Acridiens. Espèces de la faune indo-et austro-malaisienne du Museo Civico di Storia Naturale di Genova (*Ann. Mus. Genova* XXXIX 66-101).
1902. Green, *Entom. Monthly Mag.* XXXVIII 215.
1902. Bolivar, Les Orthoptères de St. Joseph's College à Trichinopoly (sud de l'Inde) (*Ann. Soc. Ent. France* LXX 580-586).
1904. Hancock, The Tettigidæ of Ceylon (*Spolia Zeylanica* II 97-157, 4 tabs.)
1906. Hancock, Wysteman's Genera Insectorum, Tetriginæ.
1907. Hancock, Studies of the Tetriginæ in the Oxford University Museum (*Trans. Entom. Soc. London*, 1907, 213-244, t. 21).
1908. Hancock, A new Ceylonese Tettigid of the genus *Eurymorphopus* (*Spol. Zeylan.* V 113-114, fig. .)
1909. Hancock, Further studies of the Tetriginæ in the Oxford University Museum (*Trans. Ent. Soc.* 1908, 387-426, t. 22).
1909. Bolivar, Nouvelles espèces d'Acridiens du Musée de Genève (Madrid, *Bol. Soc. Españ.* IX 393-408).
1909. Lefroy, Indian Insect Life, pp. 80-82.
1910. Hancock, Notes on Ceylonese Tetriginæ, with descriptions of some new species (*Spol. Zeylan.* VI 140-149, 2 figs.).
1910. Hancock, Third Paper on the Tetriginæ in the Oxford University Museum (*Trans. Ent. Soc.* 1910, 346-365, t. 49).
1910. Kirby, Synonymic Catalogue of Orthoptera, Vol. III.
1912. Hancock, Tetriginæ (Acridiinae) in the Agricul. Research Institute, Pusa, with descriptions of new species (*Ind. Agric. Ent. Mem.* IV, No. 2, pp. 131-160).
1913. Hancock, Zoological Results of the Abor Expedition, Orthoptera I (*Rec. Ind. Mus.* VIII, 311-316, t. 15).
1914. Kirby, Fauna of British India, Orthoptera, Acridiidae, pp. 11-80.
1915. Hancock, Indian Tetriginæ (*Rec. Ind. Mus.* XI 55-132, t. 14).
1918. Bolivar, Contribucion al conocimiento de la Fauna india Orthoptera (*Locustidæ vel Acridiidae*). (*Revista de la Real Academia de Ciencias Exactas, Fisicas y Naturales de Madrid*, Tomo XVI, numeros 6, 7, 8 & 9 de la segunda serie).

The following key to the sub-families of Indian Acrydidæ is adapted from Hancock (1915):—

- | | | | |
|---|---|--|-----------------|
| 1 | { | Antennæ with all the articles excepting the basal and the small atrophied apical articles deplanate triquetrous, composed of 8-10 joints; vertex transverse, very broad, or strongly acuminate produced | Tripetalocerinæ |
| | | Antennæ filiform | 2 |
| 2 | { | Anterior femora above distinctly sulcate; pronotum anteriorly produced over the head in a cornute process; antennæ with 16-22 joints | Batrachidianæ |
| | | Anterior femora more or less compressed carinate above | 3 |
| 3 | { | Frontal costa widely forked, the rami forming a frontal scutellum | Cladonotinæ |
| | | Frontal costa furcillate, but the rami diverge only moderately, or parallel, very frequently separated by a sulcus | 4 |
| 4 | { | Pronotum anteriorly truncate, or rarely angulate produced; posterior angles of the lateral lobes of pronotum turned down, more or less rounded; first hind-tarsal joint longer than third | Acrydinæ |
| | | Pronotum truncate anteriorly, posterior angles of lateral lobes laminate produced outwards, acute, or posteriorly obliquely truncate, rarely turned down | 5 |
| 5 | { | Posterior angles of lateral lobes of pronotum produced outwards, often spined; first joint of posterior tarsi longer than third; posterior tibiæ strongly ampliate, or margins dilated toward the apices; very frequently not or sparingly spinose | Scelimeninæ |
| | | Posterior angles of lateral lobes of pronotum little produced outwards, obliquely truncate behind, very rarely acute spinose; first and third joints of hind tarsi nearly equal in length | Metrodorinæ |

Keys to the genera will be found in the references quoted under each sub-family and to the species usually in the references quoted under each genus.

PUSA :
25th March, 1921.

T. BAINBRIGGE FLETCHER.

Catalogue of Indian Insects.

Part I.—ACRYDIDÆ (TETTIGIDÆ).

TRIPETALOCERINÆ.

Tripetalocerini, Kirby, F. I. Orth. p. 11
(1914) [key to genera]

Tripetalocerinæ, Hancock, Rec. Ind. Mus.
XI 55-56 (1915) [key to genera].

Tripetalocera, Westwood (1834).

Type : *ferruginea*, Westw.

Westw., Zool. Journ. V 444 (1834) ; Bol., Ann.
S. E. Belg. XXXI 190, 196, 310 (1887) ;
Hanc., Gen. Ins. Tetr. p. 4 (1906) ; Kirby,
Cat. Orth. III, 1 (1910), F. I. Orth. p. 12
(1914) [redescr.]

ferruginea, Westw., Zool. Journ. V 444, Travancore⁵
t. 22 f. 3 (1834)¹; de Haan, Temm. Malabar⁴
Verh. Orth. pp. 166, 168, t. 22 f. 11
(1842) [*Acridium (Tetrix) ferrugi-* Penang⁴
neum]²; Bol., Ann. S. E. Belg. XXXI Borneo⁴
310, t. 5 f. 28 (1887)³; Kirby, Cat. (eadem?)⁴
Orth. III 1 (1910)⁴, F. I. Orth. pp. 12-
13, ff. 15, 16 (1914)⁵.

2. Birmana, Brunner (1893).

Type : *gracilis*, Brunn.

Brunn., Ann. Mus. Genova XXXIII 113-114
(1893) ; Hanc., Gen. Ins. Tetr. p. 4 (1906) ;
Kirby, Cat. Orth. III 1 (1910), F. I. Orth.
pp. 13-14 (1914) ; Hanc., Rec. Ind. Mus.
XV 61 (1915).

gracilis, Brunn., An Mus. Genova XXXIII
114, t. 5 f. 47, (1893)¹ ; Kirby, Cat. Bhamo^{1, 4}.
Orth III¹ (1910),² F. I. Orth p. 14
(1914)³ ; Hanc., Rec. Ind. Mus. XI 61
(1915)⁴.

CLADONOTINÆ.

Cladonotini, Kirby, F. I. Orth. p. 14
(1914) [key to genera]
Cladonotinæ, Hanc., Rec. Ind. Mus. XI 56,
61 (1915) [key to genera].

3. Fieberiana, Kirby (1914).

Type : *pachymerus*, Fieb.

Plagiocephalus [*nec* Macq.], Fieber, Abh. Type : *pachymerus*, Fieb.
Königl.-böhm. Ges. Wiss. (5) III 407
(1845).

Fieberiana, Kirby, F. I. Orth. p. 14 (1914). Type : *pachymerus* Fieb.
pachymerus, Fieber, Abh. Königl.-böhm. India³
Ges. Wiss. (5) III 407 (1845) [*Plagio-*
cephalus]¹; Kirby, Cat. Orth. III 7
(1910) [*Piezotettix*]², F. I. Orth. p. 15
(1914) [redescr.]³

4. Oxyphyllum, Hancock (1909).

Type : *pennatum*, Hanc.

Hanc., T. E. S. 1908, 393 (1909), Rec. Ind.
Mus. XI 56, 61 (1915).

pennatum, Hanc., T. E. S. 1908, 393-394, Darjiling^{1, 2}
t. 22 f. 3 (1909)¹, Rec. Ind. Mus. XI
61 (1915)²

5. Deltonotus, Hancock (1904).

Type : *subcucullatus*, Wlk.

Pæcilotettix [*nec* Scudd.], Bol., Ann. S. E. Fr. Type : *gibbiceps*, Bol.
LXX 580 (1902).

Deltonotus, Hanc., Spol. Zeyl. II 107, 111 Type : *subcucullatus*,
(1904), Gen. Ins. Tetr. pp. 9, 14 (1906); Wlk.
Kirby, Cat. Orth. III 7 (1910), F. I.
Orth. p. 15 (1914) [redescr.]; Hanc.,
Rec. Ind. Mus. XI 56, 61 (1915).

gibbiceps, Bol., Ann. S. E. Fr. LXX 580 Madura^{1, 2, 3 5}
(1902) [*Pæcilotettix*]¹; Kirby, Cat. Orth.
III 7 (1910)², F. I. Orth. pp. 16-17
(1914)³; Hanc, Gen. Ins. Tetr. p. 14
(1906)⁴, Rec. Ind. Mus. XI 61 (1915)⁵.

- subcucullatus**, Wlk., Cat. Derm. Salt. Ceylon^{1, 5}
 V 830 (1871) [*Tettix subcucullata*]¹. Pundaluoya^{2, 4, 6}
Deltonotus tectiformis, Hanc., Spol. Zeyl. Hantane²
 II 107, 111, 154, t. 1 ff. 2, 2a (1904)², Kandy⁷
 Gen. Ins. Tetr. p. 14, t. 1 f. 1 (1906)³,
 T. E. S. 1907, 215 (1907)⁴.
Deltonotus subcullatus, Kirby, Cat.
 Orth. III 7 (1910)⁵, F. I. Orth. pp. 15-
 16, f. 17 (1914)⁶.
Deltonotus subcullatus, Hanc., Rec.
 Ind. Mus. XI 61 (1915)⁷.

6. *Cladonotus*, Saussure (1861).

Type: *humbertianus*, Sauss.

- Sauss., Ann. S. E. Fr. (4) I 478 (1861);
 Bolivar, Ann. S. E. Belg. XXXI 184, 192,
 208 (1887); Hancock, Spol. Zeyl. II 107,
 112-113 (1904) [redescr.], Gen. Ins. Tetrig.
 pp. 10, 16 (1906) [charact.], T. E. S. 1908
 394-395 (1909) [revised]; Kirby, Cat.
 Orth. III 9 (1910), F. I. Orth. p. 17 (1914)
 [redescr.]; Hanc., Rec. Ind. Mus. XI 56,
 61-62 (1915) [key to spp.].
humbertianus, Sauss., Ann. S. E. Fr. (4) I Ceylon^{4, 6}
 478 (1861)¹; Bol., Ann. S. E. Belg. Peradeniya^{2, 5}
 XXXI 209, t. 4 ff. 10, 10a (1887)²; Trincomali^{2, 5}.
 Hanc., Spol. Zeyl. II 107, 113-114
 (1904)³; Kirby, Cat. Orth. III 9
 (1910)⁴, F. I. Orth. pp. 17-18, f. 18
 (1914)⁵; Hanc., Rec. Ind. Mus. XI
 62 (1915)⁶.
latiramus, Hanc., Spol. Zeyl. II 107, 113, Ceylon^{3, 5}
 114-116, t. 1 ff. 1, 1a (1904)¹, Gen. Ins. Kandy^{1, 4}
 Tetrig. p. 16, t. 1 f. 3 (1906)²; Kirby,
 Cat. Orth. III 9 (1910)³, F. I. Orth.
 p. 19 (1914)⁴; Hanc., Rec. Ind. Mus.
 XI 62 (1915)⁵.
turrifer, Wlk., Cat. Derm. Salt. V 843 Ceylon¹⁻⁴
 (1871)¹; Kirby, Cat. Orth. III 9
 (1910)², F. I. Orth., pp. 18-19, f. 19

(1914)³; Hanc., Rec. Ind. Mus. XI 62
(1915)⁴.

7. Tettilobus, Hancock (1909).

Type : *spinifrons*, Hanc.

Tettilobus, Hanc., T. E. S. 1908 396 (1909), Type : *spinifrons*, Hanc
Rec. Ind. Mus. XI 56, 62 (1915).

Mnema, Kirby, F. I. Orth. p. 19 (1914). Type : *pelops*, Wlk.

pelops, Wlk., Cat. Derm. Salt. V 843 (1871) Ceylon¹⁻⁴

[*Cladonotus*]¹; Kirby, Cat. Orth. III
9 (1910) [*Cladonotus*]²; Kirby, F. I.
Orth. p. 20, ff. 20, 21 (1914) [*Mnema*]³;
Hanc., Rec. Ind. Mus. XI 62 (1915)⁴.

spinifrons, Hanc., T. E. S. 1908 396-397, Ceylon^{1, 2}
t. 22 f. 4 (1909)¹, Rec. Ind. Mus. XI
62 (1915)².

8. Gignotettix, Hancock (1909).

Type : *burri*, Hanc.

Hanc., T. E. S. 1908 397-398 (1909), Rec.
Ind. Mus. XI 56, 62 (1915).

burri, Hanc., T. E. S. 1908 398, t. 22 f. 5 Pundaluoya¹
(1909)¹, Rec. Ind. Mus. XI 62 (1915)².

9. Potua, Bolivar (1887).

Type : *coronata*, Bol. [Borneo ; Malacca].

Bol., Ann. S. E. Belg. XXXI 183, 192, 208
(1887); Hanc., Gen. Ins. Tetrig. pp. 10, 19
(1906) [charact.]; Kirby, Cat. Orth. III
11 (1910); Hanc., Rec. Ind. Mus. XI 56,
62 (1915) [charact.]

sabulosa, Hanc., Rec. Ind. Mus. XI 62-63 Satara distr.¹
(1915)¹.

SCELIMENINÆ.

Scelimenini, Kirby, F. I. Orth. pp. 20-21
(1914) [key to genera].

Scelimeninæ, Hanc., Rec. Ind. Mus. XI
56-58 [key to genera], 63 (1915).

10. *Amphibotettix*, Hancock (1906).

Type : *longipes*, Hanc. [Borneo].

Hanc., Entl. News XVIII 86 (1906), Gen. Ins. Tetrig. p. 22 (1906); Kirby, Cat. Orth. III 12 (1910); Hanc., Rec. Ind. Mus. XI 57, 63 (1915).

rosaceus, Hanc., Rec. Ind. Mus. XI 63-64 Dawna Hills¹ (1915)¹.

11. *Scelimena*, Serville (1839).

Type : *producta*, Serv. [Java].

Scelimena, Serv., Ins. Orth. p. 762 (1839); Type : *producta*, Serv. Bol., Ann. S. E. Belg. XXXI 184, 193, 215 (1887); Hanc., Spol. Zeylan. II 107, 116, 154 (1904) (redescr.), Gen. Ins. Tetrig. pp. 21, 23 (1906) [charact.]; Kirby, Cat. Orth. III 12 (1910), F. I. Orth. p. 21 (1914) [redescr.]; Hanc., Rec. Ind. Mus. XI 57, 64-65 (1915) [charact.; key to Indian spp.]

Scelymena, Sauss., Ann. S. E. Fr. (4) I 484 Type : *producta*, Serv. (1861).

Scelthymena, Bol., Ann. S. E. Fr. LXX 581 Type : *harpago*, Serv. (1902).

contracta, Wlk., Cat. Derm. Salt. V 841 India² (1871)¹; Kirby, Cat. Orth. III 13 (1910) [*Scelimena* ?]².

gavialis, Sauss., Ann. S. E. Fr. (4) I 485 Ceylon⁵ (1861) [*Scelymena*]¹.

Scelymena nodosa, Wlk., Cat. Derm. Pundaluoya^{4, 7, 9} Salt. V 840 (1871)². Dambulla^{4, 7}

Scelimena harpago [nec Serv.], Green, Ent. Maskeliya^{4, 6, 7} Mo. Mag. XXXVIII 215 (1902)³. Madulsima^{6, 8}

Scelimena gavialis, Hanc., Spol. Zeyl. Haputale⁶ II 107, 117-120, 154, t. 1 ff. 4, 4 a-c Labugama^{6, 8} (1904)⁴; Kirby, Cat. Orth. III 13 Kandy⁹ (1910)⁵; Hanc., Spol. Zeyl. VI 141-143 Pollibetta (Coorg). (1910) [habits]⁶; Kirby, F. I. Orth. pp. 24-25, f. 24 (1914)⁷; Hanc., Rec. Ind. Mus. XI 66 (1915)⁸, T. E. S. 1907, 218-219 (1907)⁹.

- harpago, Serv.**, Ins. Orthopt. p. 763 (1839) Bombay^{1, 4, 5, 7}
 [*Tetrix*]¹. Matheran⁶
Tetrix harpago, de Haan, Temm., Verh., Bandra⁷
 Orth. p. 166 (1842)². Igatpuri⁸
Scelimena harpago, Bol., Ann. S. E. Satara distr.⁸
 Belg. XXXI 216-217, t. 4 f. 13 (1887)³. Kasara⁸
Scelhymena harpago, Bol., Ann. S. E. Fr. Almora⁷
 LXX 581 (1902)⁴. Coimbatore⁷
Scelimena harpago, Kirby, Cat. Orth. S. India⁴
 III 13 (1910)⁵; Hanc., Ind. Agric. Ceylon⁴
 Ent. Mem. IV 132 (1912)⁶; Kirby,
 F. I. Orth. pp. 23-24, f. 23 (1914)⁷;
 Hanc., Rec. Ind. Mus. XI 65-66 (1915)
 [variation]⁸.
- logani, Hanc.**, Spol. Zeyl. II 107, 120-122, Ceylon³
 t. 1 ff. 5, 5 a-c (1904)¹, Gen. Ins. Tetrig. Kandy^{1, 5}
 p. 24, t. 2 f. 15 (1906)²; Kirby, Cat. Haragama^{1, 5}
 Orth. III 13 (1910)³; Hanc., Spol. Pundaluoya⁵
 Zeyl. VI. 143 (1910) [habits]⁴; Kirby, Madulsima⁴
 F. I. Orth. pp. 25-26, f. 25 (1914)⁵; Kelawewa (N. C. P.)⁷
 Hanc., Rec. Ind. Mus. XI 65 (1915)
 [charact.]⁶, T. E. S. 1907 218 (1907)⁷;
 Lefroy, Ind. Ins. Life, f. 21 on p. 76
 (1909)⁸.
- spinata, Hanc.**
- Scelimena gaviialis* [*nec* Sauss.], Kirby, Trivandrum^{1, 2}
 F. I. Orth. pp. 24-25 [part.] (1914)¹.
Scelimena spinata, Hanc., Rec. Ind. Mus.
 XI 66-67 (1915)².
- uncinata, Serv.**, Orth. pp. 763-764 (1839) Bombay^{1, 5}
 [*Tetrix*]¹.
Scelymena contracta, Wlk., Cat. Derm.
 Salt. V 851 (1871)².
Scelimena uncinata, Bol., Ann. S. E.
 Belg. XXXI 216, 218 (1887)³.
Scelimena ? uncinata, Kirby, Cat. Orth.
 III 13 (1910)⁴.
Scelimena uncinata, Kirby, F. I. Orth.
 p. 26, f. 26 (1914)⁵; Hanc., Rec. Ind.
 Mus. XI 65-66 (1915) [doubtful sp.]⁶.

12. *Eugavialidium*, Hancock (1906).

Type : *dentiumeris*, Hanc. [Borneo].

Hanc., Gen. Ins. Tetrig., pp. 21, 24 (1906);
Kirby, Cat. Orth. III 14 (1910), F. I. Orth.
p. 30 (1914) [redescr.]; Hanc., Rec. Ind.
Mus. XI 57, 67-70 (1915) [key to spp.].

angulatum, Hanc., Rec. Ind. Mus. XI 73 Calcutta¹
(1915)¹.

birmanicum, Brunn., Ann. Mus. Genova Burma^{3, 5}
XXXIII 104, t. 5 f. 37 (1893) [*Gavia- Karen Hills^{1, 4}*
lidium]¹.

Eugavialidium birmani, Hanc., Gen.
Ins. Tetrig. p. 25 (1906)².

Eugavialidium birmanicum, Kirby, Cat.
Orth. III 14 (1910)³.

Scelimena ? birmanica, Kirby, F. I. Orth.
p. 27 (1914)⁴.

Eugavialidium birmanicum, Hanc., Rec.
Ind. Mus. XI 73 (1915)⁵.

discale, Hanc.

Scelimena producta [nec Serv.], Brunn., Karen Hills^{1, 2}
Ann. Mus. Genova XXXIII 103 Sibsagar³
(1893)¹; Kirby, F. I. Orth. p. 22 [part.]
(1914)².

Eugavialidium discalis, Hanc., Rec. Ind.
Mus. XI 71 (1915)³.

feæ, Bol., R. Soc. Españ. Nat. Hist. IX Karen Hills^{1, 2}
396 (1909)¹; Hanc., Rec. Ind. Mus.
XI 73 (1915)².

flavopictum, Bol., Real. Soc. Españ. Nat. Calcutta^{1, 2}
Hist. IX 394-395 (1909)¹; Hanc., Rec.
Ind. Mus. XI 73 (1915)².

india, Hanc.

Scelimena india, Hanc., T. E. S. 1907 Cherrapunji^{1, 3}
219 (1907) [male]¹; Kirby, Cat. Orth. Assam^{1, 2, 4}
III 576 (1910)², F. I. Orth. p. 23 (1914) Shillong
[orig. descr. reprod.]³

Eugavialidium indicum, Hanc., Rec. Ind.
Mus. XI 72 (1915) [charact.]⁴

kempi, Hanc.

- Scelimena india* [nec Hanc. 1907], Hanc., Aborland^{1, 2}
 Rec. Ind. Mus. VIII 311 (1913)¹.
Eugavialidium kempi, Hanc., Rec. Ind.
 Mus. XI 72-73 (1915)².

multidentatum, Hanc.

- Scelimena producta* [nec Serv.], Kirby, Dawna Hills^{1, 2}
 F. I. Orth., p. 22 [part.] (1914)¹.
Eugavialidium multidentatum, Hanc.,
 Rec. Ind. Mus. XI 70-71 (1915)².
saussurei, Hanc., Rec. Ind. Mus. XI 74 Calcutta¹
 (1915)¹.

13. Abbasia, Kirby (1914).

Type: *subserrata*, Kirby.

- Kirby, F. I. Orth., pp. 27-28 (1914).
subserrata, Kirby, F. I. Orth. p. 28, f. 27 Tenmalai¹
 (1914)¹. [Travancore]

14. Gavialidium, Saussure (1861).

Type: *crocodilus*, Sauss.

- Sauss., Ann. S. E. Fr. (4) I 481 (1861); Bol.,
 Ann. S. E. Belg. XXXI 184, 193, 218
 (1887); Hanc., Spol. Zeyl. II 107, 122
 (1904) [redescr.], Gen. Ins. Tetrig. pp. 22,
 25 (1906) [charact.]; Kirby, Cat. Orth.
 III 14 (1910), F. I. Orth. p. 28 (1914)
 [redescr.]; Hanc., Rec. Ind. Mus. XI 57
 [charact.], 74 (1915).
alligator, Sauss., Ann. S. E. Fr. (4) I 483 Ceylon^{1, 3, 4}
 (1861) [*Scelymena*]¹; Bol., Ann. S. E. Kadugannawa⁵
 Belg. XXXI 218, 219 (1887)²; Hanc.,
 Spol. Zeyl. II 107, 122, 125 (1904)³;
 Kirby, Cat. Orth. III 15 (1910)⁴, F. I.
 Orth. pp. 29-30 (1914)⁵; Hanc., Rec.
 Ind. Mus. XI 74 (1915) [probably
 immature *crocodilus*]⁶.

- crocodilus**, Sauss., Ann. S. E. Fr. (4) I 481 Ceylon⁵
 (1861) [*Scelymena*]¹; Bol., Ann. S. E. Kadugannawa^{1, 3, 7}
 Belg. XXXI 218, 219 (1887)²; Hanc., Peradeniya⁸
 Spol. Zeyl. II 107, 122, 123-124, t. 2 Pundaluoya^{3, 7, 8, 9}
 ff. 11, 11 *a, b*. (1904)³, Gen. Ins. Tetrig. Haputale⁶
 p. 25, t. 2 ff. 16, 16*a* (1906)⁴; Kirby, Madulsima⁶
 Cat. Orth. III 14 (1910)⁵; Hanc.,
 Spol. Zeyl. VI 143-144 (1910) [habits]⁶;
 Kirby, F. I. Orth. p. 29, f. 28 (1914)⁷;
 Hanc., Rec. Ind. Mus. XI 74 (1915)⁸;
 T. E. S. 1907 220 (1907)⁹.

15. Bolotettix, Hancock (1907).

Type: *validispinus*, Hanc. [N. W. Borneo].
 Hanc., T. E. S. 1907 223 (1907); Kirby, Cat.
 Orth. III 577 (1910); Hanc., Rec. Ind.
 Mus. XI 57 [character.], 75 [key to Indian
 spp.] (1915).

- anomalus**, Hanc., Spol. Zeyl. VI 146 (1910) Madulsima¹⁻³
 [*Systolederus*]¹; Kirby, F. I. Orth. p. 46
 (1914) [*Systolederus*]²; Hanc., Rec.
 Ind. Mus. XI 76 (1915)³.
armatus, Hanc., Rec. Ind. Mus. XI 76-77 Dawna Hills¹
 (1915)¹.

exsertus, Bol.

Criotettix exsertus, Bol., Ann. S. E. Fr. Kodaikanal¹⁻⁴
 LXX 583 (1902)¹; Kirby, Cat. Orth.
 III 16 (1910)², F. I. Orth. pp. 34-35,
 f. 33 (1914)³.

Bolotettix exsertus, Hanc., Rec. Ind. Mus.
 XI 79 (1915)⁴.

- inermis**, Hanc., Rec. Ind. Mus. XI 77-78 Darjiling distr.¹ (4,000
 (1915)¹. feet)

lobatus, Hanc.,

Systolederus lobatus, Hanc., Ind. Agric. Shillong
 Ent. Mem. IV 143-144 (1912)¹. Lebong¹
Mazarredia lugubris, Kirby, F. I. Orth. Ghumti (Darjiling distr.;
 p. 55, f. 51 (1914)². 4,000 feet)³
Bolotettix lobatus, Hanc., Rec. Ind. Mus. Kurseong³
 XI 78-79 [redescr.] [1915]³.

oculatus, Bol.,

Criotettix oculatus, Bol., Ann. Mus. Kodaikanal^{2, 4, 6}
 Genova XXXIX 71 (1898)¹, Ann. Coorg⁵
 S. E. Fr. LXX 584 (1902)²; Kirby, ———
 Cat. Orth. III 16 (1910)³, F. I. Orth. Java^{3, 4, 6}
 p. 34, f. 34 on p.37 (1914)⁴; Bol., Rev. Sumatra^{3, 4, 6}
 R. Acad. Cienc. Madrid (2) XVI 283
 (1918)⁵.

Bolotettix oculatus, Hanc., Rec. Ind.
 Mus. XI 76 (1915)⁶.

pictipes, Hanc., Rec. Ind. Mus. XI 77 Shevaroy¹
 (1915).

quadratus, Hanc., Rec. Ind. Mus. XI 79-80 Singla (Darjiling distr.;
 (1915)¹. 1,500 feet)¹
 Sikkim¹

triangularis, Hanc., Rec. Ind. Mus. XI 80 Sibsagar¹
 (1915)¹.

16. Thoradonta, Hancock (1909).

Type: *dentata*, Hanc. [Penang].

Hanc., T. E. S. 1908 407 (1909), Rec. Ind.
 Mus. XI 57 [charact.], 80 (1915), T. E. S.
 1910 358-359 (1910).

apiculata, Hanc., Rec. Ind. Mus. XI 81-82 Dawna Hills¹
 (1915)¹. Tenasserim¹
 Sibsagar¹
 Singla (Darjiling distr.;
 1,500 feet)¹

sinuata, Hanc., Rec. Ind. Mus. XI 81 Satara distr.¹
 (1915)¹.

spiculoba, Hanc., Ind. Agric. Ent. Mem. IV Bihar¹
 138 (1912)¹, Rec. Ind. Mus. VIII 312-
 313 (1913) [variation]², Rec. Ind. Mus. Dibrugarh²
 XI 80-81 (1915)³. Calcutta³
 Rangoon³
 Kandy³

17. Criotettix, Bolivar (1887).

Type: *tricarinatus*, Bol.

Bol., Ann. S. E. Belg. XXXI 184, 193, 222
 (1887); Hanc., Spol. Zeyl. II 108, 128
 (1904) [redeser.], Gen. Ins. Tetrig. pp. 22,

- 27, (1906) [character.]; Kirby, Cat. Orth. III 16 (1910), F. I. Orth. p. 31 (1914) [redescri.]; Hanc., Rec. Ind. Mus. XI 58 [character.], 82-84 [key to spp.] (1915).
- æqualis**, Hanc., Ind. Agric. Ent. Mem. IV 135-136 (1912)¹, Rec. Ind. Mus. XI 87, t. 14 f. 3 (1915)². Lebong^{1, 2}
- annandalei**, Hanc., Rec. Ind. Mus. XI 87, t. 14 f. 6 (1915)¹. Paresnath (4,300 ft.)¹
- dohertyi**, Hanc., Rec. Ind. Mus. XI 86-87, t. 14 f. 4 (1915)¹. Upper Assam¹
- extremus**, Hanc., Ind. Agric. Ent. Mem. IV 132-133 (1912)¹, Rec. Ind. Mus. XI 89, t. 14 f. 11 (1915) [possibly a form of *maximus*]². Shevaroy^{1, 2}
Nilgiris (5,000 ft.)
- fastiditus**, Bol., Rev. R. Acad. Cienc. Madrid (2) XVI 284 (1918)¹. Taliparamba¹
- flavopictus**, Bol., Ann. S. E. Fr. LXX 582 (1902)¹; Kirby, Cat. Orth. III 18 (1910) [Acanthalobus]², F. I. Orth. p. 41 (1914) [Acanthalobus]³; Hanc., Rec. Ind. Mus. XI 88, t. 14 f. 10 (1915)⁴, T. E. S. 1907 220 (1907)⁵. Kodaikanal^{1, 3}
Anamalais⁴
Dawna Hills⁴
Cherrapunji⁵
- grandis**, Hanc., Ind. Agric. Ent. Mem. IV 134-135 (1912)¹, Rec. Ind. Mus. XI 89, t. 14 f. 8 (1915)². Cherrapunji^{1, 2}
Shillong
Khasis (3100 ft.)
Darjiling distr. (1,500ft.)²
Shan Hills²
- gravelyi**, Hanc., Rec. Ind. Mus. XI 88, t. 14 f. 9 (1915)¹. Ghumti (Darjiling distr.;
- indicus**, Bol., Ann. S. E. Fr. LXX 581-582 (1902)¹; Kirby, Cat. Orth. III 16 (1910)², F. I. Orth. pp. 33-34 (1914)³; Hanc., Rec. Ind. Mus. XI 85 (1915)⁴; Bol., Rev. R. Acad. Cienc. Madrid (2) XVI 283 (1918)⁵. 4,000 ft.)¹
Sikkim¹
South India¹⁻³
Burma⁶
- Criotettix maculatus*, Kirby, F. I. Orth. p. 32, f. 30 (1914)⁶.
- maximus**, Hanc., Rec. Ind. Mus. VIII 311-312, t. 15 ff. 1, 1^a (1913)¹, Rec. Ind. Mus. XI 89, t. 14 f. 7 (1915)². Aborland^{1, 2}
Janakmukh^{1, 2}
Ghumti (Darjiling distr.; 4,000 ft.)²

- montanus**, Hanc., Ind. Agric. Ent. Mem. Simla^{1, 2}
IV 133-134 (1912)¹, Rec. Ind. Mus.
XI 87, t. 14 f. 1 (1915)².
- orientalis**, Hanc., Rec. Ind. Mus. VIII 312, Dibrugarh^{1, 2}
t. 15 ff. 4, 4^a (1913)¹, Rec. Ind. Mus.
XI 85, t. 14 f. 12 (1915)².
- pallidus**, Hanc., Rec. Ind. Mus. XI 85-86, Tenasserim¹
t. 14 f. 5 (1915)¹.
- rugosus**, Bol., Ann. S. E. Belg. XXXI 228 Pegu²
(1887)¹; Brunn., Ann. Mus. Genova Rangoon³
XXXIII 105 (1893)²; Hanc., Rec. Ind. Mus. XI 84-85, t. 14 f. 2 (1915)³.
Borneo¹
- spinilobus**, Hanc., Spol. Zeyl. II 108, 129- Pundaluoya^{1, 5, 6}
131, t. 3, ff. 12, 12^{a, b} (1904)¹, Gen. Maskeliya⁴
Ins. Tetrig. p. 28, tab. f. 13 (1906)²; Travancore Coast^{6, 7}
Kirby, Cat. Orth. III 16 (1910)³; Hanc.,
Spol. Zeyl. VI 145 (1910)⁴; Kirby, F.
I. Orth. pp. 31-32 (1914)⁵; Hanc.,
Rec. Ind. Mus. XI 85, t. 14 f. 13
(1915)⁶.
- Criotettix obscurus*, Kirby, F. I. Orth.
pp. 32-33, f. 31 (1914)⁷.
- tricarinatus**, Bol., Ann. S. E. Belg. XXXI Ceylon^{1, 4}
184, 223, 224 (1887)¹, Ann. S. E. Fr. Weligama⁵
LXX 583 (1902)²; Hanc., Spol. Zeyl. Polgahawela⁵
II 108, 128-129, t. 3 ff. 15, 15^{a, b} Kadugannawa^{3, 7}
(1904)³; Kirby, Cat. Orth. III 16 Peradeniya^{3, 7, 8}
(1910)⁴; Hanc., Spol. Zeyl. VI 144 Kandy^{3, 7, 8}
(1910) [habits]⁵, Ind. Agric. Ent. Sigiriya⁸
Mem. IV 135 (1912)⁶; Kirby, F. I. Pundaluoya^{3, 7}
Orth. p. 33 f. 32 (1914)⁷; Hanc., Rec. Madulsima⁵
Ind. Mus. XI 85, t. 14 f. 14 (1915)⁸; Kodaikanal²
Bol., Rev. R. Acad. Cienc. Madrid (2) Shevaroy⁶
XVI 283 (1918)⁹. Coimbatore⁹
Virajpet (Coorg)

18. *Loxilobus*, Hancock (1904).

Type: *acutus*, Hanc.

- Hanc., Spol. Zeyl. II 108, 134 (1904), Gen.
Ins. Tetrig. pp. 22, 29 (1906) [character.];
Kirby, Cat. Orth. III 18 (1910), F. I. Orth.

- p. 41 (1914) [redescr.]; Hanc., Rec. Ind. Mus. XI 58, 91 (1915) [charact.].
- acutus, Hanc.**, Spol. Zeyl. II 108, 134—135, Pundaluoya^{1, 4}
t. 1 f. 3, t. 3 ff. 16, 16^{a, b} (1904)¹, Gen. Ceylon^{2, 3, 5}
Ins. Tetrig. p. 29, f. 14, t. 2 f. 17
(1906)²; Kirby, Cat. Orth. III 18
(1910)³, F. I. Orth. p. 42 (1914)⁴;
Hanc., Rec. Ind. Mus. XI 90 (1915)⁵,
[*nec* Hanc., Ind. Agric. Ent. Mem.
IV 137 (1912), male, = *parvispinus*.]
- assamus, Hanc.**, T. E. S. 1907 223 (1907)¹; Cherrapunji^{1, 4}
Kirby, Cat. Orth. III 577 (1910)²; Lebong³
Hanc., Ind. Agric. Ent. Mem. IV 136
(1912)³; Kirby, F. I. Orth. p. 43
(1914)⁴; Hanc., Rec. Ind. Mus. XI 90
(1915)⁵
- hancocki, Kirby,**
- Loxilobus rugosus* [*nec* Bol.], Hanc., Spol. Pundaluoya^{1, 3, 7}
Zeyl. II 108, 134, 135-136, t. 3 ff. 17, Ceylon^{2, 5}
17^{a, b} (1904)¹, Gen. Ins. Tetrig. p. 30 Weligama⁴
(1906)², T. E. S. 1908 401 (1909)³, Bassein Fort⁶
Spol. Zeyl. VI 145 (1910)⁴.
- Loxilobus hancocki*, Kirby, Cat. Orth. III
18 (1910)⁵; Hanc., Ind. Agric. Ent.
Mem. IV 137 (1912)⁶; Kirby, F. I.
Orth., pp. 42-43, f. 40 (1914)⁷; Hanc.,
Rec. Ind. Mus. XI 90 (1915)⁸.
- parvispinus, Hanc.**, Rec. Ind. Mus. XI 90- Pusa^{1, 2}
91 (1915)¹.
- Loxilobus acutus* [*nec* Hanc. 1904],
Hanc., Ind. Agric. Ent. Mem. IV
137 (1912) [male]².
- striatus, Hanc.**, Rec. Ind. Mus. XI 91 Calcutta¹
(1915)¹. Dawna Hills¹
- subulatus, Bol.,**
- Criotettix subulatus*, Bol., Ann. S. E. E. Indies¹⁻³
Belg. XXXI 185, 223, 227 (1887)¹; Ceylon⁵
Kirby, Cat. Orth. III 16 (1910)², F. I.
Orth., p. 35 (1914)³.

Loxilobus subulatus, Hanc., Rec. Ind. Mus. XI 90 (1915)⁴; Bol., Rev. R. Acad. Cienc. Madrid (2) XVI 284 (1918)⁵

vidali, Bol.,

Criotettix vidali, Bol., Ann. S. E. Belg. XXXI 185, 223, 227 (1887)¹; Brunn., Ann. Mus. Genova XXXIII 105 (1893)²; Kirby, Cat. Orth. III 16 (1910)³, F. I. Orth. pp. 35-36 (1914)⁴; Hanc., Rec. Ind. Mus. XI 89 (1915)⁵.
Loxilobus vidali, Bol., Rev. R. Acad. Cienc. Madrid (2) XVI 284 (1918)⁶.
 Karen Hills^{2, 4, 5}
 Philippines^{1, 3}

19. Acanthalobus, Hancock (1904).

Type: *miliarius*, Bol.

Hanc., Spol. Zeyl. II 108, 131 (1904), Gen. Ins. Tetrig. pp. 22, 28 (1906); Kirby, Cat. Orth. III 17 (1910), F. I. Orth. p. 36 (1914) [redescr.]; Hanc., Rec. Ind. Mus. XI 58 [character.], 91-92 [key to spp.] (1915).

bispinosus, Dalman,

Acrydium bispinosum, Dalm., Vet. Acad. Handl. 1818, p. 77¹. Burma⁸
 Karen Hills^{8, 10}
Tettix bispinosa, Dalm., Anal. Entom. p. 86 (1823)². Sibsagar¹¹
 Brahmaputra River
Tettix pallitarsis, Wlk., Cat. Derm. Salt. V 822-823 (1871)³. — above Jorhat
Tettix armigera, Wlk., Cat. Derm. Salt. V 826-827 (1871)⁴. Penang¹²
Tettix latispina, Wlk., Cat. Derm. Salt. V 837 (1871)⁵. Java^{6, 7}
 Sumatra¹⁰
Tettix bispinosus, Stål, Recens. Orth. I 150 (1873)⁶. Borneo^{7, 10}
Criotettix bispinosus, Bol., Ann. S. E. Belg. XXXI 185, 223, 226 (1887)⁷; Brunn., Ann. Mus. Genova XXXIII 105 (1893)⁸. Hongkong^{3, 4}
 China^{3, 6, 7, 9, 10}

- Acanthalobus bispinosus*, Kirby, Cat. Orth. III 17-18 (1910)⁹, F. I. Orth. pp. 37-38 (1914)¹⁰; Hanc., Rec. Ind. Mus. XI 93-94 (1915)¹¹, T. E. S. 1907, 222 (1907)¹².
- cuneatus**, Hanc., Spol. Zeyl. II 108 (1904)¹, Colombo^{2, 4}
Spol. Zeyl. II 133-134 (1904) [*A. miliarius cuneatus*]; Kirby, Cat. Orth. III 17 (1910)³, F. I. Orth. pp. 40-41, ff. 37-39 (1914)⁴; Hanc., Rec. Ind. Mus. XI 92 (1915) [apparently short-wing form of *miliarius*]⁵.
- curticornis**, Hanc., Rec. Ind. Mus. XI 92-93 (1915)¹. Satara distr.¹
- inornatus**, Wlk.,
- Tettix inornata*, Wlk., Cat. Derm. Salt. Burma^{1 5}
V 834 (1871)¹.
- Criotettix saginatus*, Bol., Ann. S. E. Belg. Rangoon^{3, 6, 7}
XXXI 185, 223, 225-226 (1887)²; Madura⁴
Brunn., Ann. Mus. Genova XXXIII
104, t. 5 f. 38 (1893)³; Bol., Ann. S. E.
Fr. LXX 583 (1902)⁴.
- Acanthalobus inornatus*, Kirby, Cat. Orth. _____
III 17 (1910)⁵, F. I. Orth. pp. 39-40,
f. 36 (1914).⁶
- Acanthalobus saginatus*, Hanc., Rec. Ind. Java^{2, 5, 6, 7A}
Mus. XI 93 (1915)⁷, T. E. S. 1907, 221 Sumatra⁵
(1907)^{7A}. China^{7A}
- Acanthalobus inornatus*, Bol., Rev. R. Acad. Cienc. Madrid (2) XVI 285
(1918) [doubts whether *inornatus* and
saginatus are really only one sp.]⁸.
- miliarius**, Bol., Ann. S. E. Belg. XXXI 185, Ceylon^{1, 4}
223, 226 (1887) [*Criotettix*]¹; Hanc., Colombo^{2, 5, 7}
Spol. Zeyl. II 108, 131, 132-133, t. 2 Galle⁵
ff. 8, 8^a (1904)², Gen. Ins. Tetrig. p. 29, Trincomali⁷
t. 2, ff. 19, 19^a (1906)³; Kirby, Cat. Kandy^{2, 6}
Orth. III 17 (1910)⁴; Hanc., Spol. Zeyl. Peradeniya^{2, 6}
VI 145 (1910)⁵; Kirby, F. I. Orth. Pundaluoya⁸
p. 37, [nec f. 34] (1914)⁶; Hanc., i ec.

Ind. Mus. XI 92 (1915)⁷, T. E. S. 1907
222 (1907)⁸.

rufescens, Kirby, F. I. Orth. pp. 38-39, Gangtok¹
f. 35 (1914).¹ Kurseong¹
Calcutta¹

20. *Tettitelum*, Hancock (1915).

Type: *hastulatum*, Kirby.

Tettitelum, Hanc., Rec. Ind. Mus. XI 94
(1915).

Tettitellum, Hanc., Rec. Ind. Mus. XI 58
(1915) [error typogr.].

hastulatum, Kirby, F. I. Orth., p. 30, Tenasserim^{1, 2}
f. 29 (1914) [*Eugavialidium*]¹.

Tettitelum hastatum, Hanc., Rec. Ind.
Mus. XI 94-95 (1915)².

METRODORINÆ.

Metrodorini, Kirby, F. I. Orth. pp. 43-44
(1914) [key to genera].

Metrodorinæ, Hanc., Rec. Ind. Mus. XI
58-59 [charact. ; key to genera], 95 (1915).

21. *Systolederus*, Bolivar (1887).

Type: *haani*, Bol. [Philippines].

Bol., Ann. S. E. Belg. XXXI 185, 194, 234
(1887); Hanc., Spol. Zeyl. II 108, 136
(1904) [redescr.], Gen. Ins. Tetrig.
pp. 30, 33 (1906) [charact.]; Kirby, Cat.
Orth. III 20 (1910), F. I. Orth. p. 44 (1914)
[redescr.]; Hanc., Rec. Ind. Mus. XI 58
[charact.], 95 (1915).

cinereus, Brunner, Ann. Mus. Genova Karen Hills.¹⁻⁴
XXXIII 105 (1893)¹; Kirby, Cat.
Orth. III 20 (1910)², F. I. Orth. pp. 44-
45, f. 41 (1914)³; Hanc., Rec. Ind.
Mus. XI 95 (1915)⁴.

greeni, Bol., Ann. S. E. Fr. LXX 584 (1902)¹; Hanc., Spol. Zeyl. II 108, 137-138, t. 2 ff. 9, 9^{a, b} (1904)², Gen. Ins. Tetrig. p. 34, t. 2 f. 14 (1906)³; Kirby, Cat. Orth. III 20 (1910)⁴; Hanc., Spol. Zeyl. VI 145-146 (1910) [habits]⁵; Kirby, F. I. Orth. p. 45, f. 42 (1914)⁶; Hanc., Rec. Ind. Mus. XI 95 (1915)⁷.
 Pundaluoya^{1, 2, 6, 7}
 Madulsima^{5, 7}
 Haputale⁵
 Maskeliya^{5, 7}
 Kadugannawa^{2, 6}
 Kodaikanal^{1, 6}
 Nilgiris⁷
 Virajpet (Coorg)
 Pusa

22. *Hancockia*, Kirby (1914).

Type : *portentosa*, Kirby.

Kirby, F. I. Orth. p. 46 (1914).

portentosa, Kirby, F. I. Orth. pp. 46-47, f. 43 (1914)¹, Travancore¹

23. *Eurymorphopus*, Hancock (1906).

Type : *cunctatus*, Bol. [New Caledonia].

Hanc., Gen. Ins. Tetrig. pp. 30, 35 (1906); Kirby, Cat. Orth. III 21 (1910), F. I. Orth. pp. 48-49 (1914) (redescr.); Hanc., Rec. Ind. Mus. XI 58 [character.], 95 (1915).

latilobus, Hanc., Spol. Zeyl. V 113-114, f. 1 (1908)¹; Kirby, F. I. Orth. p. 49, ff. 46, 47 (1910)²; Hanc., Rec. Ind. Mus. XI 95 (1915)³.
 Undugoda¹⁻³
 Kandy³

24. *Spadotettix*, Hancock (1910)

Type : *fletcheri*, Hanc.

Hanc., Spol. Zeyl. VI 146-147 (1910), Ind. Agric. Ent. Mem. IV 141-142 (1912); Kirby, F. I. Orth. p. 70 (1914) [redescr.]; Hanc., Rec. Ind. Mus. XI 58 [character.], 95 (1915).

fletcheri, Hanc., Spol. Zeyl. VI 147-148, figs. (1910)¹; Kirby, F. I. Orth. p. 71 (1914)²; Hanc., Rec. Ind. Mus. XI 95 (1915)³.
 Madulsima^{1, 2}

provertex, Hanc., Ind. Agric. Ent. Mem. Shevaroy's¹
 IV 142-143 (1912)¹, Rec. Ind. Mus.
 XI 95 (1915)².

25. Apterotettix, Hancock (1904).

Type : *obtusus*, Hanc.

Hanc., Spol. Zeyl. II 108, 140 (1904), Gen.
 Ins. Tetrig. pp. 30, 35 (1906) [character.];
 Kirby, Cat. Orth. III 21 (1910), F. I. Orth.
 p. 47 (1914) [redescr.]; Hanc., Rec. Ind.
 Mus. XI 58 [character.], 95 (1915).

obtusus, Hanc., Spol. Zeyl. II 108, 140-
 141, t. 3 ff. 13, 13^{a, b} (1904)¹, Gen. Ins. Pundaluoya^{1, 4}
 Tetrig. p. 35, f. 16 (1906)²; Kirby, Peradeniya^{1, 4, 5}
 Cat. Orth. III 21 (1910)³, F. I. Orth.
 p. 48, ff. 44, 45 (1914)⁴; Hanc., Rec.
 Ind. Mus. XI 96 (1915)⁵.

26. Amphinotus, Hancock (1915).

Type : *pygmæus*, Hanc.

Hanc., Rec. Ind. Mus. XI 58-59, 96
 (1915).

pygmæus, Hanc., Rec. Ind. Mus. XI 96-
 97 (1915)¹. Hakgala¹
 Pundaluoya¹

27. Mazarredia, Bolivar (1887).

Type : *gemella*, Bol. [Philippines].

Bol., Ann. S. E. Belg. XXXI 185, 194, 236-
 237 (1887); Brunn., Ann. Mus. Genova
 XXXIII 106 (1893); Hanc., Spol. Zeyl.
 II 108, 138 (1904) [redescr.], Gen. Ins.
 Tetrig. pp. 32, 42 (1906) [character.]; Kirby,
 Cat. Orth. III 24 (1910), F. I. Orth. pp. 48-
 51 (1914) [redescr.]; Hanc., Rec. Ind. Mus.
 XI 59 [character.], 97-99 [key to spp.] (1915).

convergens, Brunn., Ann. Mus. Genova Karen Hills^{1, 3}
 XXXIII 106, 107-108, t. 5 f. 40 Tenasserim^{1, 3}
 (1893)¹; Kirby, Cat. Orth. III 26
 (1910)², F. I. Orth. p. 52 (1914)³;
 Hanc., Rec. Ind. Mus. XI 101 (1915)⁴.

- crisulata**, **Bol.**, Ann. S. E. Fr. LXX 584-585 (1902)¹; Kirby, Cat. Orth. III 25 (1910)², F. I. Orth. pp. 52-53 (1914)³; Hanc., Rec. Ind. Mus. XI 102 (1915)⁴. Madura^{1, 3}
- dubia**, **Hanc.**, Ind. Agric. Ent. Mem. IV 141 (1912), Rec. Ind. Mus. XI 101 (1915)². Lebong^{1, 2}
Cherrapunji
- ghumtiana**, **Hanc.**, Rec. Ind. Mus. XI 100-101 (1915)¹. Ghumti (Darjiling distr.)¹
- inæqualis**, **Brunn.**, Ann. Mus. Genova XXXIII 106-107, t. 5 f. 39 (1893)¹; Kirby, Cat. Orth. III 25 (1910)², F. I. Orth. p. 53, f. 49 (1914)³; Hanc., Rec. Ind. Mus. XI 99 (1915)⁴. Karen Hills^{1, 3}
- insignis**, **Kirby**, F. I. Orth. p. 54, f. 50 (1914)¹. Bhim Tal¹
Shillong
Coorg
- insularis**, **Bol.**, Ann. S. E. Belg. XXXI 237, 239 (1887)¹; Hanc., Spol. Zeyl. II 108, 138-140, t. 2 ff. 7, 7^{a, b} (1904)², T. E. S. 1907 230 (1907)³; Kirby, Cat. Orth. III 25 (1910)⁴, F. I. Orth. p. 52 (1914)⁵; Hanc., Rec. Ind. Mus. XI 102 (1915)⁶. Ceylon¹
Pundaluoya^{2, 3, 5}
Hantana^{2, 5}
Peradeniya⁵
Kandy³
- laticeps**, **Bol.**, Bol. Soc. Españ. IX 399 (1909)¹; Hanc., Rec. Ind. Mus. XI 102 (1915)². Upper Assam^{1, 2}
- latifrons**, **Hanc.**, Ind. Agric. Ent. Mem. IV 139 (1912)¹, Rec. Ind. Mus. XI 102 (1915)². Lebong^{1, 2}
- lativertex**, **Brunn.**, Ann. Mus. Genova XXXIII 106, 108, t. 5 f. 41 (1893)¹; Kirby, Cat. Orth. III 26 (1910)², F. I. Orth. p. 54 (1914)³; Hanc., Rec. Ind. Mus. XI 100 (1915)⁴. Karen Hills^{1, 3}
- ophthalmica**, **Bol.**, Bol. Soc. Españ. IX 399 (1909)¹; Hanc., Rec. Ind. Mus. XI 102 (1915)². Sibsagar^{1, 2}
- perplexa**, **Hanc.**, Rec. Ind. Mus. XI 101 (1915)¹. Sikkim¹

- sculpta**, Bol., Ann. S. E. Belg. XXXI 237, "East Indies"¹
 238 (1887)¹; Brunn., Ann. Mus. Genova Tenasserim^{2, 4, 5}
 XXXIII 107 (1893)²; Hanc., T. E. S. Karen Hills^{2, 5}
 1908 405 (1909)³; Kirby, Cat. Orth. Pegu^{2, 4, 5}
 III 25 (1910)⁴, F. I. Orth. p. 51 [*nec* Assam³
 f. 48] (1914)⁵; Hanc., Rec. Ind. Mus. Sukna⁶
 XI 100 (1915)⁶; Boliv., Rev. R. Acad.
 Cienc. Madrid (2) XVI 285 (1918)⁷.
- sikkimensis**, Bol., Bol. Soc. Españ. IX Sikkim²
 398-399 (1909)¹; Hanc., Rec. Ind.
 Mus. XI 100 (1915)².
- singlaënsis**, Hanc., Rec. Ind. Mus. XI Singla (Darjiling distr.;
 99-100 (1915)¹. 1,500 feet)¹

28. *Xistra*, Bolivar (1887).

Type: *gogorzæ*, Bol. [Philippines].

- Bol., Ann. S. E. Belg. XXXI 186, 194, 242
 (1887); Brunn., Ann. Mus. Genova
 XXXIII 108 (1893); Hanc., Gen. Ins.
 Tetrig. pp. 32, 45 (1906) [charact.]; Kirby,
 Cat. Orth. III 27 (1910), F. I. Orth. p. 55
 (1914) [redescr.]; Hanc., Rec. Ind. Mus.
 XI 59 [charact.], 102 (1915).

dubia, Brunn.,

- Xistra? dubia*, Brunn., Ann. Mus. Karen Hills^{1, 3}
 Genova XXXIII 108-109, t. 5 f. 42
 (1893)¹; Kirby, Cat. Orth. III 27
 (1910)², F. I. Orth. p. 56 (1914)³.
Xistra dubia, Hanc., Rec. Ind. Mus. XI
 102 (1915)⁴.

- sikkimensis**, Hanc., Rec. Ind. Mus. XI Sikkim¹
 102-103 (1915)¹.

- stylata**, Hanc., T. E. S. 1907 231-232 Puttalam^{1, 4}
 (1907)¹, T. E. S. 1908 409 (1909)²; Kandy²
 Kirby, Cat. Orth. III 577 (1910)³, Nagody (S. Kanara)⁶
 F. I. Orth. p. 56 (1914)⁴; Hanc., Rec. Coorg
 Ind. Mus. XI 102 (1915)⁵; Boliv., Rev.
 R. Acad. Cienc. Madrid (2) XVI 286
 (1918)⁶.

29. *Xistrella*, Bolivar (1909).Type: *dromadaria*, Bol.

Bol., Bol. Soc. Españ. IX 400-401 (1909);
Hanc., Rec. Ind. Mus. XI 59 [character.],
103 (1915).

dromadaria, Bol., Bol. Soc. Españ. IX Sikkim^{1, 2}
401 (1909)¹; Hanc., Rec. Ind. Mus.
XI 103 (1915) [female descr.]²

30. *Lamellitettix*, Hancock (1904).Type: *acutus*, Hanc.

Hanc., Spol. Zeyl. II 107, 125-126 (1904),
Gen. Ins. Tetrig. pp. 31, 41 (1906)
[character.]; Kirby, Cat. Orth. III 24
(1910), F. I. Orth. p. 49 (1914) [redescr.];
Hanc., Rec. Ind. Mus. XI 59 [character.],
103 (1915).

acutus, Hanc., Spol. Zeyl. II 107, 126-128, Maskeliya^{1, 5}
t. 2 fl. 6, 6^{a, b} (1904)¹, Gen. Ins. Tetrig. Howaheta³
p. 42, t. 2 fl. 21, 21^a (1906)², T. E. S.
1908 404 (1909)³; Kirby, Cat. Orth.
III 24 (1910)⁴, F. I. Orth. p. 50
(1914)⁵.

fletcheri, Hanc., Rec. Ind. Mus. XI 103 Anamalais¹
(1915)¹.

pluricarinatus, Hanc., T. E. S. 1908 404 Deltota^{1, 3}
(1909)¹, Spol. Zeyl. VI 148 (1910)², Maskeliya²
Rec. Ind. Mus. XI 103 (1915)³.

31. *Hyboella*, Hancock (1915).

Hanc., Rec. Ind. Mus. XI 59 [character.],
104-109 [Key to spp.] (1915).

Type: *tentata*, Hanc.

acuteterminatus, Brunn., Ann. Mus. Pegu^{1, 3}
Genova XXXIII 112 (1893) [*Copto-*
tettix]¹; Kirby, Cat. Orth. III 55
(1910)², F. I. Orth. pp. 75-76, f. 70
(1914)³ [*Coptotettix*]; Hanc., Rec. Ind.
Mus. XI 106 (1915)⁴.

- angulifrons**, Hanc., Rec. Ind. Mus. XI 108-109 (1915)¹. Dawna Hills¹
- conioptica**, Hanc., Rec. Ind. Mus. XI 109 (1915)¹. Singla (Darjiling distr.; 1,500 feet)¹
Assam¹
- dilatata**, de Haan,;
- Acridium (Tetrix) dilatatum*, de Haan, Temm., Verh. Orth. pp. 167, 169, t. 22 f. 12 (1842)¹. Karen Hills^{3, 5}
- Tetrix dilatatus*, Bol., Ann. S. E. Belg. XXXI 187, 257, 260 (1887)²; Brunn., Ann. Mus. Genova XXXIII 109 (1893)³. Java^{1, 2, 4, 5}
- Acrydium dilatatum*, Kirby, Cat. Orth. III 45 (1910)⁴, F. I. Orth. p. 68 (1914)⁵.
- Hyboella dilatata*, Hanc., Rec. Ind. Mus. XI 107 (1915)⁶.
- latifrons**, Brunn.,
- Coptotettix latifrons*, Brunn., Ann. Mus. Genova XXXIII 112, t. 5 f. 44 (1893)¹; Kirby, Cat. Orth. III 55 (1910)², F. I. Orth. pp. 74-75, ff. 67-69 (1914)³. Pegu^{1, 3}
Pusa³
- Hyboella latifrons*, Hanc., Rec. Ind. Mus. XI 106 (1915)⁴.
- nullipennis**, Hanc., Rec. Ind. Mus. VIII 314, t. 15 ff. 2, 2^a (1913) [*Coptotettix*]¹, (N. E. Assam)
Rec. Ind. Mus. XI 106 (1915)². Janakmukh^{1, 2}
- obesa**, Hanc., Rec. Ind. Mus. XI 107-108 (1915)¹. Ghumti (Darjiling distr.; 4,000 feet)¹
- problematica**, Bol.,
- Coptotettix problematicus*, Bol., Real Soc. Españ. IX 401 (1909)¹. Upper Assam^{1, 2}
- Hyboella problematica*, Hanc., Rec. Ind. Mus. XI 109 (1915)².
- tentata**, Hanc., Rec. Ind. Mus. XI 106-107 (1915)¹. Sibsagar¹

tumida, Hanc.,

Coptotettix tumidus, Hanc., Rec. Ind. Dibrugarh^{1, 2}

Mus. VIII 313-314, t. 15 ff. 3, 3^a
(1913)¹.

Hyboella tumida, Hanc., Rec. Ind. Mus.
XI 106 (1915)².

ACRYDINÆ.

Acrydiini, Kirby, F. I. Orth. p. 57 (1914)
[key to genera].

Tettiginæ, Hanc., Rec. Ind. Mus. XI 59-60
[charact. ; key to genera], 109 (1915).

32. Teredorus, Hancock (1906).

Type: *stenofrons*, Hanc. [Peru]

Teredorus, Hanc., Gen. Ins. Tetrigr. pp. 51-52
(1906); Kirby, Cat. Orth. III 30
(1910); Hanc., Rec. Ind. Mus. XI
109-110 (1915).

Terederus [error typogr.], Hanc., Rec. Ind.
Mus. XI 59 [charact.] (1915).

carmichaeli, Hanc., Rec. Ind. Mus. XI Singla (Darjiling distr.;
110 (1915)¹. 1,500 feet)¹

frontalis, Hanc., Rec. Ind. Mus. XI 110 Simla Hills (5,000 ft.)¹
(1915)¹.

33. Paratettix, Bolivar (1887).

Type: *meridionalis*, Rambur [Mediterranean
Region].

Batrachidia [nec Serv.], Sauss., Rev. Zool. (2)
XIII 401 (1861).

Paratettix, Bol., Ann. S. E. Belg. XXXI 187,
195, 270 (1887); Finot, Faune France,
Orth. pp. 100, 169 (1889); Lugger,
Orthopt. Minnesota pp. 196, 199 (1897);
Morse, Biol. Centr. Am., Orth. II 5, 11
(1900); Hanc., Tettig. N. Amer. pp. 36,
39, 108 (1902); Jacobs. and Bianchi,
Prem. i Lozhn. Ross. Imp. pp. 164, 175, 211

- (1902); Blatchley, Rep. Indiana Dep. Geology XXVII 218, 227 (1903); Hanc., Spol. Zeyl. II 108, 110, 144 (1904) [redescr.], Gen. Ins. Tetrig. pp. 52, 55 (1906) [character.]; Kirby, Cat. Orth. III 32 (1910); Hanc., Ind. Agric. Ent. Mem. IV 149 (1912); Kirby, F. I. Orth. pp. 60-61 (1914) [redescr.]; Hanc., Rec. Ind. Mus. XI 60 [character.], 111-112 [key to spp.] (1915).
- alatus**, Hanc., Rec. Ind. Mus. XI 113-114 (1915)¹. Ghumti (Darjiling distr.; 4,000 feet)¹
- curtipennis**, Hanc., Ind. Agric. Ent. Mem. IV 146-147 (1912) [*Coptotettix*]¹, Rec. Ind. Mus. XI 113 (1915)². Lebong¹
Darjiling distr. (4,000 feet)²
Upper Shillong (6,000 feet)
- dorsifer**, Wlk.
Tettix dorsifera, Wlk., Cat. Derm. Salt. V 825 (1871)¹. Bombay¹⁻³
Acrydium dorsiferum, Kirby, Cat. Orth. III 45 (1910)².
Paratettix? *dorsifer*, Kirby, F. I. Orth. p. 63, f. 55 (1914)³.
- hirsutus**, Brunn., Ann. Mus. Genova XXXIII 110, t. 5 ff. 43^{a, b} (1893)¹; Kirby, Cat. Orth. III 33 (1910)², F. I. Orth. p. 65, f. 57 (1914)³; Hanc., Rec. Ind. Mus. XI 112 (1915)⁴. Teinzo^{1, 3}
Katha^{1, 3}
Bhamo^{1, 3}
Sibsagar⁴
- latipennis**, Hanc., Rec. Ind. Mus. XI 113 (1915)¹. Moradabad¹
- rotundatus**, Hanc., Rec. Ind. Mus. XI 112-113 (1915)¹. Tezpur¹
- scaber**, Thunberg
Acrydium scabrum, Thnb., Nova Acta Upsal, VII 159 (1815)¹. South India^{5, 8}
Coimbatore¹⁰
Tettix subpustulata, Wlk., Cat. Derm. Salt. V 819 (1871)². Palamcottah¹⁰
Kezanathum¹⁰
Tettix scaber, Stål, Recens. Orth. I 149 (1873)³. Bangalore¹⁰
Nandyal¹⁰

- Paratettix scaber*, Bol., Ann. S. E. Belg. Annampallee¹⁰
 XXXI 188, 279 (1887)⁴, Ann. S. E. ———
 Fr. LXX 585-586 (1902)⁵. South Africa^{2, 3, 7}
- Paratettix scaber ugandensis*, Karny, Sitz. Zanzibar^{4, 8}
 Akad. Wiss. Wien. CXVI 26 (1907)⁶. East Africa⁸
- Paratettix scaber*, Kirby, Cat. Orth. III Gaboon^{4, 8}
 32 (1910)⁷, F. I. Orth. p. 62 (1914)⁸; Aldabra⁹
 Bol., Tr. Linn. Soc. (2) XV 267 (1912)⁹, Seychelles⁹
 Rev. R. Acad. Cienc. Madrid (2) XVI
 287 (1918)¹⁰.
- semihirsutus**, Brunn., Ann. Mus. Genova. Karen Hills^{1, 3}
 XXXIII 110-111 (1893)¹; Kirby, Cat.
 Orth. III 33 (1910)², F. I. Orth. p. 66,
 f. 58 (1914)³; Hanc., Rec. Ind. Mus.
 XI 113 (1915)⁴.

34. *Acrydium*, Geoffroy (1762).

Type: *subulatum*, Linn. [Europe, W. and N.
 Asia]

- Acrydium*, Geoff., Hist. Ins. I 390 (1762); Type: *subulatum*, Linn.
 Fabr., Syst. Entom. p. 278 (1775);
 Rossi, Faun. Etrusc. p. 262 (1790);
 Oliv., Enc. Méthod. Ins. VI 209
 (1791); Leach, Edinb. Encycl. IX
 120 (1815); Curt., Brit. Ent. X t. 439
 (1833); Steph., Ill. Brit. Ent., Mandib.
 VII 32 (1835); Kirby, Cat. Orth. III
 35 (1910), F. I. Orth. pp. 66-67 (1914);
 Hanc., Rec. Ind. Mus. XI 60 [character],
 114 (1915).
- Bulla*, Schrank, Enum. Ins. Austr. p. 242 Type:?
 (1781).
- Acridium*, Schrank, Fauna Boica II 30 Type:?.....
 (1801); Zett., Orth. Suec. p. 104
 (1821), Faun. Ins. Lapp. I 452 (1828);
 Phil., Orthopt. Berol. p. 40 (1830);
 Zett., Ins. Lapp. p. 250 (1840).
- Tetrix*, Latr., Hist. Nat. Crust. Ins. III 284 Type: *subulata*, Linn.
 (1802), XII 161 (1804); St. Farg. and
 Serv., Encycl. Meth. Ins. X 598
 (1825); Serv., Ann. Sci. Nat. XXII

264, 291 (1831), *Ins. Orthopt.* p. 757 (1839); Wesm., *Bull. Acad. Brux.* V 596 (1838); Burm., *Handb. Ent.* II 657 (1838); Borck, *Skand. Rätv. Ins.* p. 138 (1848); Finot, *Orth. France* pp. 39, 82 (1883), *Faune France Orth.* pp. 99, 165 (1889); Jacobs. and Bianchi, *Prem. i Lozhn. Ross. Imp.* pp. 174, 208 (1902); Hanc., *Tett. N. Amer.* pp. 35, 38, 66, 68 (1902), *Spol. Zeyl.* II 108, 110, 141 (1904) [redescr.], *Gen. Ins. Orth.* pp. 52, 57 (1906) [character.], *Ind. Agric. Ent. Mem.* IV 149 (1912); Lucas, *Brit. Orth.* p. 206 (1920).

Tettix, Charp., *Germ. Zeitschr. Ent.* III 315 (1841); Fieb., *Abh. Böhn. Ges.* (5) III 408 (1844); Fisch. *Waldh., Orth. Russ.* p. 347 (1846); Fisch., *Orth. Eur.* p. 215 (1853); Fieber, *Lotos* III 139 (1853), *Syn. Eur. Orth.* p. 29 (1854); Türk, *Wien. Ent. Mon.* VI 206 (1862); Friv., *Ertek. Termész. Köreb.* I (12) p. 188 (1868); Stâl, *Recens. Orth.* I 146 (1873); Bol., *Ortopt. Esp.* p. 161 (1876), *An. Soc. Esp.* V 367 (1876); Brunn., *Prodr. Eur. Orth.* pp. 87, 234 (1882); Bol., *Ann. S. E. Belg.* XXXI 187, 195, 257 (1887), *Ann. Sci. Nat. Porto* IV 221, V 35 (1897-98); Burr, *Brit. Orth.* pp. 33, 46 (1897); Auriv., *Ent. Tidskr.* XXI 241, 247 (1900); Tümpel, *Geradfl. Mitteleuropas* pp. 220, 253 (1900); Hanc., *Spol. Zeyl.* II 141-142 (1904) [redescr.].

Type: *subulatus*, Linn.

bipunctatum, Linn.

Gryllus Bulla bipunctatus, Linn., *Syst. Nat.* (ed. X) p. 427 (1758)¹, *Sulz., Gesch. Ins.* p. 82, t. 8 f. 6 (1776)².
 Kodaikanal¹⁴, ¹⁷
 Sikkim¹⁸

- Acrydium scutellatum*, de Geer, Mém.
Ins. III 483, t. 23 f. 15 (1773)³.
- Acrydium bipunctatum*, Herbst, Fuess.
Arch. p. 189, t. 52 f. 1 (1786)⁴.
- Acrydium opacum*, Herbst, Fuess. Arch. —————
p. 190, t. 52 f. 2 (1786)⁵.
- Acrydium gibbum*, Oliv., Enc. Meth. Ins. Persian Gulf¹⁸
VI 233 (1791)⁶.
- Tetrix bipunctata*, Serv., Ins. Orthopt. N. & W. Asia¹⁶
p. 765 (1839)⁷.
- Tettix schrankii*, Fieb., Abhandl. Böhm. N. Africa¹⁶
Ges. (5) III 412, t. 10 ff. 17-19 (1843)⁸.
- Tettix bipunctatus*, Brunn., Prodr. Eur. Europe^{16, 18, 19}
Orth. pp. 234, 235 (1882)⁹; Bol., Ann. S. E. Belg. XXXI 263 (1887)¹⁰; Burr.
Brit. Orth. p. 46, t. 4 f. 2 (1897)¹¹;
Tümpel, Geradfl. Mitteleuropas pp.
220, 253, t. 20 (1900)¹².
- Tetrix bipunctata*, Jacobs. and Bianchi,
Prem. i Lozhn. Ross. Imp. pp. 174,
209, t. 7 (1902)¹³.
- Tettix bipunctatus*, Bol., Ann. S. E. Fr.
LXX 585 (1902)¹⁴.
- Tetrix bipunctata*, Hanc., T. E. S. 1910
359 (1910)¹⁵.
- Acrydium bipunctatum*, Kirby, Cat. Orth.
III 39-42 (1910) [numerous other refs.
and synonyms]¹⁶, F. I. Orth. pp. 68-
69, ff. 59, 60 (1914)¹⁷; Hanc., Rec. Ind.
Mus. XI 114 (1915)¹⁸.
- Tetrix bipunctatus*, Lucas Brit. Orth.
pp. 212-217, t. 21 f. 1 (1920)¹⁹

ceylonum, Hanc.

- Tettix atypicalis ceylonus*, Hanc, Spol. Pundaluoya^{1, 2, 4}
Zeyl. II 108, 142, 143-144 (1904)¹. Maskeliya^{1, 4}
- Tetrix atypicalis ceylonus*, Hanc., T. E. S. Peradeniya⁵
1907 239 (1907)².
- Acrydium ceylonicum*, Kirby, Cat. Orth.
III 45 (1910)³.
- Acrydium atypicale ceylonum*, Kirby, F.
I. Orth. pp. 67-68 (1914)⁴.

Acrydium ceylonicum, Hanc., Rec. Ind.
Mus. XI 115 (1915) [short-wing form
of *variegatum*, Bol.]⁵.

cingalense, Wlk.

- Tettix cingalensis*, Wlk., Cat. Derm. Ceylon^{1, 2, 8, 12, 11}
Salt. V 827 (1871)¹. Madulsima¹⁰
- Paratettix variegatus*, Bol., Ann. S. E. Kandy^{4, 13}
Belg. XXXI 188, 272, 280 (1887)²; Peradeniya^{4, 13, 19}
Hanc., Spol. Zeyl. II 108, 144-145 Pundaluoya⁵
(1904)³. Dikoya^{4, 13}
- Tettix atypicalis*, Hanc., Spol. Zeyl. II - Ratnapura¹⁴
108, 142-143, t. 3 ff. 14, 14 a, b(1904)⁴. Pondichery¹¹
- Tettix atypicalis*, Hanc., T. E. S. 1907 Taliparamba¹⁵
239 (1907)⁵. (Malabar)
- Tettix variegatus*, Hanc., T. E. S. 1908 Singla (Darjiling distr.;
413 (1909)⁶. 1,500 feet)¹⁴
- Euparatettix cingalensis*, Kirby, Cat. Moulmein¹⁴
Orth. III 31 (1910)⁷.
- Paratettix variegatus*, Kirby, Cat. Orth. Java⁶
III 33 (1910)⁸.
- Acrydium atypicalis*, Kirby, Cat. Orth.
III 45 (1910)⁹.
- Euparatettix variegatus*, Hanc., Spol.
Zeyl. VI 148 (1910)¹⁰.
- Tettix variegatus*, Hanc., Ind. Agric.
Ent. Mem. IV 149 (1912)¹¹.
- Paratettix cingalensis*, Kirby, F. I. Orth.
pp. 63-64, f. 56 (1914)¹².
- Acrydium atypicale*, Kirby, F. I. Orth.
p. 67 (1914)¹³.
- Acrydium variegatum*, Hanc., Rec. Ind.
Mus. XI 114-115 (1915)¹⁴.
- Paratettix cingalensis*, Bol., Rev. R.
Acad. Cienc. Madrid (2) XVI 287
(1918)¹⁵.

discale, Wlk.

- Tettix discalis*, Wlk., Cat. Derm. Salt. V Ceylon^{1, 2}
826 (1871)¹.
- Acrydium discalis*, Kirby, Cat. Orth. III
45 (1910)².

indicum, Bol.

- Paratettix indicus*, Bol., Ann. S. E. Belg. "East Indies"¹
 XXXI 188, 272, 281 (1887)¹, Ann. S. E. South India^{2, 5}
 Fr. LXX 585 (1902)²; Kirby, Cat. Coimbatore⁶
 Orth. III 33 (1910)³; Hanc., Ind. Bihar^{4, 7}
 Agric. Ent. Mem. IV 150 (1912)⁴; —————
 Kirby, F. I. Orth. pp. 64-65 (1914)⁵; China^{1-3, 5}
 Bol., Rev. R. Acad. Cienc. Madrid (2)
 XVI 288 (1916)⁶.
Acrydium indicum, Hanc., Rec. Ind.
 Mus. XI 115 (1915)⁷.

indum, Olivier.

- Acrydium indum*, Oliv., Enc. Méth. Ins. "East Indies"^{1, 2}
 VI 234 (1791)¹.
Acrydium ? indum, Kirby, Cat. Orth.
 III 49 (1910)².
Acrydium indicum, Kirby, F. I. Orth.
 p. 77 (1914)³; Bol., Rev. R. Acad.
 Cienc. Madrid (2) XVI 288 (1918)⁴.

lineosum, Wlk.

- Tettix lineosa*, Wlk., Cat. Derm. Salt. V, Bombay^{1, 2}
 Suppl. p. 90 (1871)¹.
Acrydium lineosum, Kirby, Cat. Orth.
 III 45 (1910)².

mundum, Wlk., Cat. Derm. Salt. V 823- North India^{1, 2}
 824 (1871) [*Tettix munda*]¹; Kirby, Cat.
 Orth. III 45 (1910)²; Hanc., Rec. Ind.
 Mus. XI 115 (1915)³.

nigricolle, Wlk., Cat. Derm. Salt. V 826 Bombay^{1, 2}
 (1871) [*Tettix nigricollis*]¹; Kirby, Cat.
 Orth. III 45 (1910)².

obliquiferum, Wlk., Cat. Derm. Salt. V Bombay^{1, 2}
 825 (1871) [*Tettix obliquifera*]¹; Kirby,
 Cat. Orth. III 45 (1910)².

quadriplagiatum, Wlk.

- Tettix quadriplagiata*, Wlk., Cat. Derm. North India¹⁻³
 Salt. V 828 (1871)¹.

- Hedotettix quadriplagiata*, Kirby, Cat. Orth. III 49 (1910)².
- Acrydium quadriplagiatum*, Kirby, F. I. Orth. p. 69, f. 61 (1914)³.
- tectitergum**, Hanc., Rec. Ind. Mus. XI 115-116 (1915)¹ Hoshangabad¹
Surat¹.
- umbriferum**, Wlk., Cat. Derm. Salt. V 824 (1871) [*Tettix umbrifera*]¹; Kirby, Cat. Orth. III 45 (1910)² Bombay^{1, 2}
- vittiferum**, Wlk., Cat. Derm. Salt. V 824-825 (1871) [*Tettix vittifera*]¹; Kirby, Cat. Orth. III 45 (1910)² Bombay^{1, 2}
- 35. Coptotettix, Bolivar (1887).**
- Type: *asperatus*, Bol. [Cochin China].
- Bol., Ann. S. E. Belg. XXXI 188, 195, 287 (1887); Hanc., Spol. Zeyl. II 108, 111, 152 (1904) [redescr.], Gen. Ins. Tetrig. pp. 52, 65 (1906) [charact.]; Kirby, Cat. Orth. III 55 (1910), F. I. Orth. p. 74 (1914) [redescr.]; Hanc., Rec. Ind. Mus. XI 60 [charact.], 116-118 [key to spp.] (1915).
- annandalei**, Hanc., Rec. Ind. Mus. XI 119 (1915)¹ Singla (Darjiling distr.; 1,500 feet)¹
Calcutta¹
N. Shan Hills¹
- artolobus**, Hanc., Rec. Ind. Mus. XI 121 (1915)¹ Ceylon¹
- capitatus**, Bol., Ann. S. E. Belg. XXXI 188, 287, 289 (1887)¹; Brunna., Ann. Mus. Genova XXXIII 111 (1893)²; Kirby, Cat. Orth. III 55 (1910)³, F. I. Orth. pp. 76-77 (1914)⁴; Hanc., Rec. Ind. Mus. XI 118 (1915)⁵; Bol., Rev. R. Acad. Cienc. Madrid (2) XVI 288 (1918)⁶ Karen Hills ²⁻⁴
Somwarpet (Coorg)⁶
Java^{1, 3}
- conspersus**, Hanc., Rec. Ind. Mus. XI 119-120 (1915)¹ Sibsagar ¹
Siliguri¹
Bhim Tal¹

- fossulatus**, Bol., Ann. S. E. Belg. XXXI 188, 287, 288-289 (1887)¹; Hanc., Spol. Zeyl. II 108, 153 (1904)²; Kirby, Cat. Orth. III 55 (1910)³, F. I. Orth. p. 76-(1914)⁴; Hanc., Rec. Ind. Mus. XI 118 (1915)⁵. Ceylon¹⁻⁴
Puri distr.⁵
- indicus**, Hanc., Ind. Agric. Ent. Mem. IV 144-145 (1912)¹, Rec. Ind. Mus. VIII 313 (1913) [female descr.]², Rec. Ind. Mus. XI 120 (1915)³. Kalyan¹
Satara distr.³
Aborland³
- interruptus**, Bol., Ann. S. E. Belg. XXXI 189, 288, 291-292 (1887)¹; Brunn., Ann. Mus. Genova XXXIII 113 (1893)²; Kirby, Cat. Orth. III 56 (1910)³, F. I. Orth. p. 78 (1914)⁴; Hanc., Rec. Ind. Mus. XI 118 (1915)⁵. Rangoon^{2, 4}
Bhamo^{2, 4}
Karen Hills^{2, 4}
Java^{1, 3, 4}
- parvulus**, Hanc., Ind. Agric. Ent. Mem. IV 145-146 (1912)¹, Rec. Ind. Mus. VIII 314 (1913)², Rec. Ind. Mus. XI 118 (1915)³. Chapra^{1, 3}
Calcutta³
Kushtea (Bengal)³
Singla (Darjiling distr.; 1,500 feet)³
Sikkim³
Janakmukh²
Kobo²
Dibrugarh²
- retractus**, Hanc., Rec. Ind. Mus. XI 120-121 (1915)¹. Pusa¹
- testaceus**, Bol., Ann. S. E. Belg. XXXI 189, 288, 291 (1887)¹; Hanc., Spol. Zeyl. II 108, 153-154 (1904)²; Kirby, Cat. Orth. III 56 (1910)³, F. I. Orth. p. 77 (1914)⁴; Hanc., Rec. Ind. Mus. XI 121 (1915)⁵. Ceylon¹⁻⁴
Chota Nagpur⁵

36. *Hedotettix*, Bolivar (1887).

Type: *gracilis*, de Haan.

- Bol., Ann. S. E. Belg. XXXI 188, 195, 283 (1887); Hanc., Spol. Zeyl. II 108, 111, 148 (1904) [redescr.], Gen. Ins. Tetrig. pp. 52, 60 (1906) [character.]; Kirby, Cat. Orth. III 49 (1910), F. I. Orth. p. 71 (1914)

[redescr.]; Hanc., Rec. Ind. Mus. XI 60
[charact.], 121-122 [key to spp.] (1915).

- attenuatus**, Hanc., Spol. Zeyl. II 108, Colombo^{1, 4, 5}
149, 151-152, t. 3 ff. 18, 18 *a, b* (1904)¹, Trincomali⁵
Gen. Ins. Tetrig. p. 60, f. 23 (1906)²; Puri⁵
Kirby, Cat. Orth. III 50 (1910)³, F. I. Sibsagar⁵
Orth. p. 73 (1914)⁴; Hanc., Rec. Ind. Assam-Bhutan Frontier⁵
Mus. XI 123 (1915)⁵.
- crstitergus**, Hanc., Rec. Ind. Mus. XI Hoshangabad¹
122 (1915)¹.
- costatus**, Hanc., Ind. Agric. Ent. Mem. Bengal^{1, 2}
IV 147-148 (1912)¹, Rec. Ind. Mus. Bihar^{1, 2}
XI 123 (1915)². Nepal²
Allahabad²

gracilis, de Haan.

- Acridium (Tetrix) gracile*, de Haan, Throughout India (to
Temm., Verh., Orth. pp. 167, 169 Peshawar), Burma and
(1842)¹. Ceylon.
- Hedotettix gracilis*, Bol., Ann. S. E. Belg. XXXI 188, 283, 284 (1887)².
- Heditettix festivus*, Bol., Ann. S. E. Belg. Siam¹¹
XXXI 188, 284, 286, t. 5 ff. 24, 24a Sumatra¹¹
(1887)³. Java^{2, 4, 8, 11, 15}
Celebes^{11, 15}
- Hedotettix gracilis*, Brunn., Ann. Mus. Genova XXXIII 111 (1893)⁴; Bol.,
Ann. S. E. Fr. LXX 586 (1902) [variation]⁵; Hanc., Spol. Zeyl. II 108, 149-
151, t. 3 ff. 19, 19 *a, b*. (1904)⁶.
- Hedotettix gracilis abortus*, Hanc., Spol. Zeyl. II 108, 149, 151 (1904)⁷.
- Hedotettix gracilis*, Hanc., T. E. S. 1907 240 (1907)⁸, T. E. S. 1908 422 (1909)⁹.
- Hedotettix gracilis abortus*, Hanc., T. E. S. 1908 422 (1909)¹⁰.
- Hedotettix gracilis*, Kirby, Cat. Orth. III 49-50 (1910)¹¹; Hanc., Spol. Zeyl. VI 149 (1910) [habits]¹², Ind. Agric. Ent. Mem. IV 147 (1912)¹³.
- Hedotettix diminutus*, Hanc., Ind. Agric. Ent. Mem. IV 148-149 (1912)¹⁴.

Hedotettix gracilis, Kirby, F. I. Orth.
p. 72, ff. 64, 65 (1914)¹⁵; Hanc., Rec.
Ind. Mus. XI 123 (1915)¹⁶; Bol., Rev.
R. Acad. Cienc. Madrid (2) XVI
288 (1918)¹⁷.

grossus, Hanc., Rec. Ind. Mus. XI 124 Singla (Darjiling distr.;
(1915)¹. 1,500 feet)¹

lineifer, Wlk.

Tettix lineifera, Wlk., Cat. Derm. Salt. V Bombay¹⁻³
824 (1871)¹.

Acrydium lineiferum, Kirby, Cat. Orth.
III 45 (1910)².

Hedotettix? *lineiferum*, Kirby, F. I.
Orth. p. 73, f. 66 (1914)³.

37. Euparatettix, Hancock (1904).

Type : *personatus*, Bol.

Euparatettix, Hanc., Spol. Zeyl. II 108, 111, Type : *personatus*, Bol.
145 (1904), Gen. Ins. Tetrigr. pp. 51,
55 (1906) [character.]; Kirby, Cat.
Orth. III 31 (1910); Hanc., Ind.
Agric. Ent. Mem. IV 150-151 (1912)
[key to spp.]; Kirby, F. I. Orth. p. 57
(1914) [redescr.]; Hanc., Rec. Ind.
Mus. XI 60 [character.], 124-125 [key
to spp.] (1915).

Ergatettix, Kirby, F. I. Orth. pp. 69-70 (1914). Type : *scabripes*, Bol.
(=*tarsalis*, Kby.)

balteatus, Wlk.

Tettix balteata, Wlk., Cat. Derm. Salt. V South India¹⁻³
828-829 (1871)¹. Salem distr.⁴

Acrydium balteatum, Kirby, Cat. Orth. Palur⁴
III 45 (1910)².

Paratettix balteatus, Kirby, F. I. Orth.
pp. 61-62, f. 54 (1914)³.

Euparatettix balteatus, Bol., Rev. R.
Acad. Cienc. Madrid (2) XVI 286
(1918)⁴.

corpulentus, Hanc., Ind. Agric. Ent. Throughout India (to
Mem. IV 153 (1912), Rec. Ind. Mus. Peshawar), Burma,
XI 126 (1915)². and Ceylon.

personatus, Bol.

Paratettix personatus, Bol., Ann. S. E. Throughout India (to
Belg. XXXI 188, 278 (1887)¹; Brunn., Peshawar), Burma,
Ann. Mus. Genova XXXIII 109 and Ceylon.
(1893)².

Euparatettix personatus, Hanc., Spol. ———
Zeyl. II 108, 145, 146-148, t. 2 ff. 10, Java^{1, 3, 5, 6, 10}
10 a, b, t. 3 ff. 20, 20a, b. (1904)³, Gen. Philippines^{1, 3, 10}
Ins. Tetrig. p. 55, t. 3 f. 32 (1906)⁴,
T. E. S. 1907 238 (1907)⁵, T. E. S.
1908 409 (1909)⁶, Spol. Zeyl. VI 148-
149 (1910) [habits]⁷; Kirby, Cat. Orth.
III 31 (1910)⁸; Hanc., Ind. Agric. Ent.
Mem. IV 152 (1912)⁹; Kirby, F. I. Orth
p. 58, ff. 52, 53 (1914)¹⁰; Hanc., Rec.
Ind. Mus. XI 125 (1915)¹¹.

personatus birmanicus, Hanc., Rec. Rangoon¹
Ind. Mus. XI 125-126 (1915)¹. Assam-Bhutan Frontier.¹

scabripes, Bol.

Paratettix scabripes, Bol., Ann. Mus. South India²
Genova XXXIX 76 (1898)¹, Ann. Tinnevelly⁶
S. E. Fr. LXX 585 (1902)²; Kirby, Salem distr.⁶
Cat. Orth. III 35 (1910)³. Coimbatore⁶
Ergatettix tarsalis, Kirby, F. I. Orth. Taliparamba (Malabar)⁶
p. 70, ff. 62, 63 (1914)⁴. Calcutta⁴
Euparatettix scabripes, Kirby, F. I. Rajshahi (Bengal)⁵
Orth. p. 59 (1914)⁵; Bol., Rev. R. Acad.
Cienc. Madrid (2) XVI 286 (1918)⁶. ———
Java^{2, 3}
Sumatra^{2, 5}

tenuis, Hanc., Ind. Agric. Ent. Mem. IV Throughout United
151-152 (1912)¹, Rec. Ind. Mus. VIII Prov., Bihar, Bengal
315 (1913)², Rec. Ind. Mus. XI 126 and Assam.
(1915)³. Dawna Hills³

variabilis, Bol.,

Paratettix variabilis, Bol., Ann. S. E. Anuradhapura⁷
Belg. XXXI 188, 271, 276-277 (1887)¹, Madura^{1, 4, 5}
Ann. S. E. Fr. LXX 585 (1902); Puri⁷

- Hanc., T. E. S. 1907 236 (1907)³; Kirby, Cat. Orth. III 33 (1910)⁴, F. I. Orth. pp. 62-63 (1914)⁵. Asansol⁷
Berhampur⁷
Rajshahi⁷
- Euparatettix variabilis*, Hanc., Ind. Agric. Ent. Mem. IV 150 (1912)⁶, Rec. Ind. Mus. XI 126 (1915)⁷. Rajmahal⁷
Pusa
Shillong
Mandalay⁷
-
- Borneo^{1, 3, 5}
Philippines^{1, 4, 5}
- 38. Indatettix, Hancock (1915).**
- Type : *nodulosus*, Hanc.
- Hanc., Rec. Ind. Mus. XI 60 [charact.], 127-129 [key to spp.] (1915).
- callosus, Hanc.,** Rec. Ind. Mus. XI 130 (1915)¹. Singla (Darjiling distr. ; 1,500 feet.)¹
- crassipes, Hanc.,** Ind. Agric. Ent. Mem. IV 153-164 (1912) [*Euparatettix*]¹, Rec. Ind. Mus. XI 130-131 (1915)². Throughout United Prov., Bihar, Bengal Gauhati.
- crassipes bengalensis, Hanc.,** Ind. Agric. Ent. Mem. IV 154-155 (1912) [*Euparatettix bengalensis*]¹, Rec. Ind. Mus. XI 131 (1915)². Pusa¹
Muzaffarpur¹
Dacca¹
- crassipes hybridus, Hanc.,** Rec. Ind. Mus. XI 131 (1915)¹. Dibrugarh¹
Madhupur¹
Nepal¹
- crassipes var. A, Hanc.,** Rec. Ind. Mus. XI 131 (1915)¹. Chapra¹
Pusa¹
- interruptus, Brunn.,**
- Paratettix interruptus*, Brunn., Ann. Mus. Genova XXXIII 109-110 (1893)¹. Karen Hills^{1, 4}
Bhamo^{1, 4}
- Euparatettix interruptus*, Hanc., T. E. S. 1910 359-360 (1910)²; Kirby, Cat. Orth. III 31 (1910)³, F. I. Orth. p. 59 [? f. 48 on p. 51] (1914)⁴; Bol., Rev. R. Acad. Cienc. Madrid (2) XVI 59 (1918)⁵. Teinzo^{1, 4}
Damukdia⁶
Madhupur⁶
Calcutta⁶
Puri⁶
Coimbatore⁵

- Indatettix interruptus*, Hanc., Rec. Ind. Mus. XI 130 (1915)⁶. Marikuppam⁶
Travancore⁶
Batticaloa (Ceylon)⁶
Johore²
- interruptus* var. **A**, Hanc., Rec. Ind. Mus. XI 130 (1915)¹. Satara distr.¹
- interruptus lobulatus*, Hanc., Rec. Ind. Mus. XI 130 (1915)¹. Hoshangabad¹
- nodulosus*, Hanc., Ind. Agric. Ent. Mem. IV 155-156 (1912) [*Euparatettix*]¹, Rec. Ind. Mus. XI 129 (1915)². Throughout United Prov., Bihar and Bengal.
Garhwal²
Nagpur
Satara²
- parvus*, Hanc.
- Euparatettix parvus*, Hanc., Spol. Zeyl. II 108, 145-146 (1914)¹. Elephant Pass (Ceylon)^{1, 3}
Weligama (Ceylon)³
- Euparatettix pilosus*, Hanc., T. E. S. 1908 409-410 (1909)², Spol. Zeyl. VI 149 (1910)³. S. Malabar
Mysore²
Central India²
- Euparatettix parvus*, Kirby, Cat. Orth. III 31 (1910)⁴; Hanc., Ind. Agric. Ent. Mem. IV 156 (1912)⁵, Rec. Ind. Mus. VIII 315 (1913)⁶; Kirby, F. I. Orth. p. 60 (1914)⁷. Puri⁸
Calcutta⁸
Chaumahani (Barisal)
Igatpuri⁸
Amangarh (U. P.)⁸
- Indatettix parvus*, Hanc., Rec. Ind. Mus. XI 129 (1915)⁸. Dhampur (U. P.)⁸
Bhim Tal⁸
Peshawar
Dibrugarh⁶
Aborland⁶
Chapra⁵
Pusa⁵

BATRACHIDIANÆ.

- Batrachidiini*, Kirby, F. I. Orth. p. 78 (1914).
- Batrachidinae*, Hanc., Rec. Ind. Mus. XI 60 [charact.], 131 (1915).

39. *Saussurella*, Bolivar (1887).

Type: *cucullifera*, Wlk. (*cornuta*, Bol., *nec* de Haan.)

Bol., Ann. S. E. Belg., XXXI 189, 196, 303 (1887); Hanc., Gen. Ins. Tetrig. pp. 67, 71 (1906); Kirby, Cat. Orth. III 61 (1910); Hanc., Ind. Agric. Ent. Mem. IV 156-157 [key to spp.] (1912); Kirby, F. I. Orth. pp. 78-79 (1914) [redescr.]; Hanc., Rec. Ind. Mus. XI 60 [character.], 131 (1915).

cucullifera, -Wlk.

Tettix cucullifera, Wlk., Cat. Derm. Salt. Pegu^{3, 4, 6, 7}
V 829 (1871)¹. Sibsagar⁸

Saussurella cornuta [*nec* de Haan], Bol.,
Ann. S. E. Belg. XXXI 189, 303 Amoy^{1, 5, 6}
(1887)²; Brunn., Ann. Mus. Genova Hongkong⁶
XXXIII 113, t. 5 f. 45 (1893)³; Kirby,
Cat. Orth. III 61 (1910) [part.]⁴

Saussurella cucullifera, Kirby, Cat.
Orth. III 61 (1910)⁵, F. I. Orth. p. 79,
f. 71 (1914)⁶.

Saussurella brunneri, Hanc., Ind.
Agric. Ent. Mem. IV 156 (1912)⁷, Rec.
Ind. Mus. XI 131-132 (1915)⁸.

curticornu, Hanc., Ind. Agric. Ent. Mem. Pusa^{1, 2}
IV 158-159 (1912)¹, Rec. Ind. Mus.
XI 131 (1915)².

decurva, Brunn., Ann. Mus. Genova Pegu^{1, 3}
XXXIII 113, t. 5 f. 46 (1893)¹; Kirby, Tenasserim⁴
Cat. Orth. III 61 (1910)², F. I. Orth. N. Lakhimpur⁴
pp. 79-80 (1914)³; Hanc., Rec. Ind.
Mus. XI 131 (1915)⁴.

indica, Hanc., Ind. Agric. Ent. Mem. IV Lebong^{1, 2}
157-158 (1912)¹, Rec. Ind. Mus. XI
131 (1915)².

TOTAL 169 species.

INDEX TO GENERA.

(NOTE.—The numbers refer to the numbers of the genera and not of the pages.)

- | | |
|----------------------------------|------------------------------------|
| Abbasia, 13 | Indatettix, 38 |
| Acanthalobus, 19 | Lamellitettix, 30 |
| <i>Acridium</i> (Acrydium) | Loxilobus, 18 |
| Acrydium, 34 | Mazarredia, 27 |
| Amphibotettix, 10 | <i>Mnema</i> (Tettilobus) |
| Amphinotus, 26 | Oxyphyllum, 4 |
| Apterotettix, 25 | Paratettix, 33 |
| <i>Batrachidia</i> (Paratettix) | <i>Plagiocephalus</i> (Fieberiana) |
| Birmana, 2 | Potua, 9 |
| Bolotettix, 15 | Saussurella, 39 |
| <i>Bulla</i> (Acrydium) | <i>Scelhymena</i> (Scelimena) |
| Cladonotus, 6 | Scelimena, 11 |
| Coptotettix, 35 | <i>Scelymena</i> (Scelimena) |
| Criotettix, 17 | Spadotettix, 24 |
| Deltonotus, 5 | Systolederus, 21 |
| <i>Ergatettix</i> (Euparatettix) | Teredorus, 32 |
| Eugavialidium, 12 | <i>Tetrix</i> (Acrydium) |
| Euparatettix, 37 | Tettilobus, 7 |
| Eurymorphosus, 23 | Tettitelum, 20 |
| Fieberiana, 3 | <i>Tettix</i> (Acrydium) |
| Gavialidium, 14 | Thoradonta, 16 |
| Gignotettix, 8 | Tripetalocera, 1 |
| Hancockia, 22 | Xistra, 28 |
| Hedotettix, 36 | Xistrella, 29 |
| Hyboella, 31 | |

INDEX TO SPECIES.

(NOTE.—Names *in italics* are treated as synonyms and should be looked up under the names given in roman letters. The generic name noted against each specific name is that of the genus in which it is placed in this Catalogue. The species in each genus are placed in alphabetical order and the position of the genus is shown in the Index to Genera.)

- abortus*, Hedotettix gracilis
acuteterminatus, Hyboella
acutus, Lamellitettix
 „ Loxilobus
æqualis, Criotettix
alatus, Paratettix
alligator, Gavialidium
angulatum, Eugavialidium
angulifrons, Hyboella
annandalei, Coptotettix
 „ Criotettix
anomalus, Bolotettix
apiculata, Thoradonta
armatus, Bolotettix
armigera, Acantholobus bispinosus
artolobus, Coptotettix
assamus, Loxilobus
attenuatus, Hedotettix
atypicalis, *Tettix* (*Acrydium cingalense*)
- balteatus*, Euparatettix
bengalensis, Indatettix crassipes
bipunctatum, Acrydium
birmanicum, Eugavialidium
birmanicus, Euparatettix personatus
bispinosus, Acantholobus
brunneri, Saussurella cucullifera
burri, Gignotettix
- callosus*, Indatettix
capitatus, Coptotettix
carmichaeli, Tereorus
ceylonicum, Acrydium ceylonum
ceylonum, Acrydium
cinereus, Systolederus
cingalense, Acrydium
conioptica, Hyboella
conspersus, Coptotettix
contracta, Scelimena
contracta, Scelimena uncinata
convergens, Mazarredia
cornuta, Saussurella cucullifera
corpulentus, Euparatettix
costatus, Hedotettix
crassipes, Indatettix
crstitergus, Hedotettix
cratulata, Mazarredia
crocodilus, Gavialidium
cucullifera, Saussurella
cuneatus, Acantholobus
- curticornis*, Acantholobus
curticornu, Saussurella
curtipennis, Paratettix
- decurva*, Saussurella
dilatata, Hyboella
diminutus, Hedotettix gracilis
discale, Acrydium
 „ Eugavialidium
dohertyi, Criotettix
dorsifer, Paratettix
dromadaria, Xistrella
dubia, Mazarredia
 „ Xistra
- exsertus*, Bolotettix
extremus, Criotettix
- fastiditus*, Criotettix
fea, Eugavialidium
ferruginea, Tripetalocera
festivus, Hedotettix gracilis
flavopictum, Eugavialidium
flavopictus, Criotettix
fletcheri, Lamellitettix
 „ Spadotettix
fossulatus, Coptotettix
frontalis, Teradorus
- gavialis*, Scelimena
ghumtiana, Mazarredia
gibbiceps, Deltonotus
gibbum, Acrydium bipunctatum
gracilis, Birmana
 „ Hedotettix
grandis, Criotettix
graveli, Criotettix
greeni, Systolederus
grossus, Hedotettix
- hancocki*, Loxilobus
harpago, Scelimena
hastulatum, Tettitelum
hirsutus, Paratettix
humbertianus, Cladonotus
hybridus, Indatettix crassipes
- inæqualis*, Mazarredia
india, Eugavialidium
indica, Saussurella

- indicum*, Acrydium
indicus, Coptotettix
 „, Criotettix
indum, Acrydium
inermis, Bolotettix
inornatus, Acanthalobus
insignis, Mazarredia
insularis, Mazarredia
interruptus, Coptotettix
 „, Indatettix

kempi, Eugavialidium

laticeps, Mazarredia
latifrons, Hyboella
 „, Mazarredia
latilobus, Eurymorphopus
latipennis, Paratettix
latiramus, Cladonotus
latispina, Acanthalobus bispinosus
lativertex, Mazarredia
lineifer, Hedotettix
lineosum, Acrydium
lobatus, Bolotettix
lobulatus, Indatettix interruptus
logani, Scelimena
lugubris, Mazarredia (Bolotettix lobatus)

maculatus, Criotettix indicus
maximus, Criotettix
miliarius, Acanthalobus
montanus, Criotettix
multidentatum, Eugavialidium
mundum, Acrydium

nigricolle, Acrydium
nodosa, Scelimena gavialis
nodulosus, Indatettix
nullipennis, Hyboella
obesa, Hyboella
obliquiferum, Acrydium
obscurus, Criotettix spinilobus
optus, Apterotettix
oculatus, Bolotettix
opacum, Acrydium bipunctatum
ophthalmica, Mazarredia
orientalis, Criotettix

pachymerus, Fieberiana
pallidus, Criotettix
pallitarsis, Acanthalobus bispinosus
parvispinus, Loxilobus
parvulus, Coptotettix
parvus, Indatettix
pelops, Tettilobus
pennatum, Oxyphyllum
perplexa, Mazarredia
personatus, Euparatettix
pictipes, Bolotettix
pilosus, Indatettix parvus

pluricarinatus, Lamellitettix
portentosa, Hancockia
problematica, Hyboella
provertex, Spadotettix
pygmæus, Amphinotus

quadratus, Bolotettix
quadriplagiatum, Acrydium

retractus, Coptotettix
rosaceus, Amphibotettix
rotundatus, Paratettix
rufescens, Acanthalobus
rugosus, Criotettix
rugosus, Loxilobus hancocki

sabulosa, Potua
saginata, Acanthalobus inornatus
saussurei, Eugavialidium
scaber, Paratettix
scabripes, Euparatettix
schranksii, Acrydium bipunctatum
sculpta, Mazarredia
scutellatum, Acrydium bipunctatum
semihirsutus, Paratettix
sikkimensis, Mazarredia
 „, Xistra
singlaensis, Mazarredia
sinuata, Thoradonta
spiculoba, Thoradonta
spinata, Scelimena
spinifrons, Tettilobus
spinilobus, Criotettix
striatus, Loxilobus
stylata, Xistra
subcucullatus, Deltonotus
subpustulata, Paratettix scaber
subserrata, Abbasia
subulatus, Loxilobus

tarsalis, Euparatettix scabripes
tectiformis, Deltonotus subcucullatus
tectitergum, Acrydium
tentata, Hyboella
tenuis, Euparatettix
testaceus, Coptotettix
triangularis, Bolotettix
tricarinatus, Criotettix
tumida, Hyboella
turrifer, Cladonotus

ugandensis, Paratettix scaber
umbriferum, Acrydium
uncinata, Scelimena

variabilis, Euparatettix
variegatus, Paratettix (Acrydium cingalense)
vidali, Loxilobus
vittiferum, Acrydium